

DIHEDRAL GALOIS COVERS OF ALGEBRAIC VARIETIES AND THE SIMPLE CASES

FABRIZIO CATANESE, FABIO PERRONI

Dedicated to Ugo Bruzzo on the occasion of his 60-th birthday.

ABSTRACT. In this article we investigate the algebra and geometry of dihedral covers of smooth algebraic varieties. To this aim we first describe the Weil divisors and the Picard group of divisorial sheaves on normal double covers. Then we provide a structure theorem for dihedral covers, that is, given a smooth variety Y , we describe the algebraic “building data” on Y which are equivalent to the existence of such covers $\pi: X \rightarrow Y$. We introduce then two special very explicit classes of dihedral covers: the simple and the almost simple dihedral covers, and we determine their basic invariants. For the simple dihedral covers we also determine their natural deformations. In the last section we give an application to fundamental groups.

CONTENTS

1. Introduction	2
2. Direct images of sheaves under Galois covers	4
2.1. The non-Galois case	9
3. Line bundles and divisorial sheaves on flat double covers	10
4. Line bundles on hyperelliptic curves	13
4.1. Powers of divisorial sheaves on double covers	17
5. Dihedral field extensions and generalities on dihedral covers	18
5.1. Dihedral field extensions	19
5.2. Structure of D_n -covers	21
5.3. D_3 -covers and triple covers.	30
6. Simple and almost simple dihedral covers and their invariants	32
6.1. Invariants of simple dihedral covers	35
6.2. Almost simple dihedral covers	36
7. Deformations of simple dihedral covers	40

Keywords: Galois covers of algebraic varieties; classification theory of algebraic varieties.

The present work took place in the framework of the DFG Forschergruppe 790 ‘Classification of algebraic surfaces and compact complex manifolds’ and of the ERC-2013-Advanced Grant - 340258- TADMICAMT. The second author was also supported by the grant FRA 2015 of the University of Trieste, PRIN “Geometria delle Varietà Algebriche” and INDAM.

8. Examples and applications	41
8.1. Simple dihedral covers of projective spaces	41
8.2. An application to fundamental groups	43
References	44

1. INTRODUCTION

A main issue in the classification theory of algebraic varieties is the construction of interesting and illuminating examples. For instance in the book of Enriques [Enr49] one can see that a recurrent method is the one of considering the minimal resolution S of double covers $f : X \rightarrow Y$; these are called ‘piani doppi’, double planes, when $Y = \mathbb{P}^2$.

Burniat [Bur66] considered more general bidouble covers, i.e., Galois covers $f : X \rightarrow Y$ with group $G = (\mathbb{Z}/2)^2$, and some work of the first author [Cat84], [Cat98] was focused on looking at the invariants and the deformations of bidouble covers, deriving basic results for the moduli spaces of surfaces.

Comessatti [Com30] was the first to study Galois coverings with abelian group G , and their relations to topology, while Pardini [Par91] described neatly the algebraic structure of such coverings, their invariants and deformations.

Just to give a flavour of the result: when G is a cyclic group of order n and Y is factorial, normal G -coverings correspond to building data (L, D_1, \dots, D_{n-1}) consisting of reduced effective divisors D_1, \dots, D_{n-1} without common components, and of the isomorphism class of a divisor L such that $nL \equiv \sum_i iD_i$ (\equiv is the classical notation for linear equivalence). The theorem is the scheme counterpart of the field theoretic description

$$\mathbb{C}(X) = \mathbb{C}(Y)[z]/(z^n - \prod_i \delta_i^i).$$

Here $D_i = \{\delta_i = 0\}$ and the branch locus \mathcal{B}_f is the union of the divisors D_i .

In Pardini’s theorem one can replace the field \mathbb{C} by any algebraically closed field of characteristic coprime to n , but over the complex numbers we have a more general result, namely, the extension of the Riemann existence theorem due to Grauert and Remmert: normal schemes with a finite covering $f : X \rightarrow Y$, and with branch locus contained in a divisor \mathcal{B} correspond to conjugacy classes of monodromy homomorphisms $\mu : \pi_1(Y \setminus \mathcal{B}) \rightarrow \mathfrak{S}_n$. The scheme is a variety (i.e., irreducible) iff $\text{Im}(\mu)$ is a transitive subgroup, and Galois if it is simply transitive (similarly for Abelian coverings in [BaCa08] the criterion for irreducibility of X was explicitly given).

On the other hand, the goal of finding explicit algebraic equations for the case of Galois coverings with non Abelian group G is like the quest for the Holy Graal, and in this paper, widely extending previous results

of Tokunaga [Tok94], we give a full characterization of Galois coverings $\pi: X \rightarrow Y$ with Y smooth and with Galois group the dihedral group D_n of order $2n$.

The underlying idea is of course the same and is easy to explain in general terms: we can factor $\pi: X \rightarrow Y$ as the composition of a cyclic covering of order n , $p: X \rightarrow Z := X/H$, (here $H \subset D_n$ is the group of rotations) and a (singular) double covering $q: Z \rightarrow Y$. A main new technical tool, developed in sections 3 and 4, consists in describing Weil divisors and the Picard group of divisorial sheaves on a normal double cover Z . To make the results more appealing, we describe in detail the very special case of the Picard group of hyperelliptic curves, especially the algebraic determination of torsion line bundles; this is an extension of a previous partial result by Mumford [Mum84], it boils down to determinantal equations for triples of polynomials in one variable, and bears interesting similarities with the resultant of two polynomials.

In section 5 we describe the general theorem, whose application in concrete cases is however not so straightforward. For this reason we concentrate in the next sections on two special very explicit classes of dihedral covers of algebraic varieties: the simple and the almost simple dihedral covers.

The underlying idea of simple covers is the one of giving a schematic equation which looks exactly as the equation describing the field extension. The first well known instance is the one of a simple cyclic cover: X is given by an equation

$$z^n = F,$$

where z is a fibre variable on the line bundle \mathbb{L} associated to a divisor L on Y , and F is a section in $H^0(\mathcal{O}_Y(nL))$.

In [Cat89] the analysis of everywhere nonreduced moduli spaces was based also on the technical notion of an almost simple cyclic cover, given as the locus X inside the \mathbb{P}^1 -bundle $\mathbb{P}(\mathbb{L}_0 \oplus \mathbb{L}_\infty)$, associated to a pair of two line bundles $\mathbb{L}_0, \mathbb{L}_\infty$, defined by the equation

$$z_0^n a_\infty = z_\infty^n a_0, \quad a_0 \in H^0(\mathcal{O}_Y(nL_0)), \quad a_\infty \in H^0(\mathcal{O}_Y(nL_\infty)),$$

and where the divisors $E_0 = \{a_0 = 0\}$ and $E_\infty = \{a_\infty = 0\}$ are disjoint (else the covering is not finite).

Let us now explicitly describe the equations of a simple dihedral covering: X is contained in a rank two split vector bundle of the form $\mathbb{L} \oplus \mathbb{L}$ over Y , and is defined by equations

$$\begin{cases} u^n + v^n &= 2a, & a \in H^0(\mathcal{O}_Y(nL)), \\ uv &= F, & F \in H^0(\mathcal{O}_Y(2L)). \end{cases}$$

The dihedral action is generated by an element σ of order n such that

$$u \mapsto \zeta u, v \mapsto \zeta^{-1}v,$$

where ζ is a primitive n -th root of 1, and by an involution τ exchanging u with v .

The branch locus is here the divisor $\mathcal{B} = \{a^2 - F^n = 0\}$, Z is defined by $z^2 = a^2 - F^n$, and the cyclic covering X is defined by $u^n = z + a$: it is smooth if the two divisors $\{a = 0\}$ and $\{F = 0\}$ intersect transversally and \mathcal{B} is smooth outside of $F = 0$.

In particular, as it is well known from the work of Zariski [Zar29] and many other followers, the fundamental group $\pi_1(Y \setminus \mathcal{B})$ is non Abelian and admits a surjection to the group D_n (irreducibility is granted if there are points where $\{a = 0\}$ and $\{F = 0\}$, since there the covering is totally ramified).

In the case of simple dihedral coverings, the covering $X \rightarrow Z$ is only ramified in the A_{n-1} singularities of Z , the points where $z = a = F = 0$. In order to get a covering with more ramification we introduce then the almost simple dihedral covers, defined this time on the fibre product of two \mathbb{P}^1 -bundles. X is the subset of $\mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \times_Y \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L})$ defined by the following two equations:

$$\begin{cases} u_1 v_1 - u_0 v_0 F & = 0 \\ a_\infty v_1^n u_0^n - 2a_0 v_0^n u_0^n + a_\infty v_0^n u_1^n & = 0, \end{cases}$$

where, v_1, u_1 are fibre coordinates on \mathbb{L} , v_0, u_0 are fibre coordinates on $\underline{\mathbb{C}}$ (so that $([u_0 : u_1], [v_0 : v_1])$ are projective fibre coordinates on $\mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \times_Y \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L})$).

As we said earlier, we want to find the invariants and study the deformations of the varieties that we construct in this way. We first do this in full generality for any Galois covering in section 2; later we write more explicit formulae for simple and almost simple dihedral coverings. We apply these in the case of easy examples, deferring the study of more complicated cases to the future.

One interesting application, to real forms of curves with cyclic symmetry, and related loci in the moduli space of real curves, shall be given in a forthcoming joint work with Michael Lönne [CLP].

Finally, in this paper we consider also the non Galois case: namely, coverings of degree n whose Galois group is D_n . In doing so, we establish connections with the theory of triple coverings of algebraic varieties described by Miranda [Mir85] (if the triple covering is not Galois, then its Galois group is $D_3!$).

2. DIRECT IMAGES OF SHEAVES UNDER GALOIS COVERS

We consider (algebraic) varieties which are defined over the field of complex numbers \mathbb{C} , though many results remain valid for complex manifolds and for algebraic varieties defined over an algebraically closed field of characteristic p not dividing $2n$.

Definition 2.1. Let Y be a variety and let G be a finite group. A **Galois cover** of Y with group G , shortly a **G -cover** of Y , is a finite morphism $\pi: X \rightarrow Y$, where X is a variety with an effective action by G , such that π is G -invariant and induces an isomorphism $X/G \cong Y$.

A **dihedral cover** of Y is a D_n -cover of Y , where D_n is the dihedral group of order $2n$.

If $\pi: X \rightarrow Y$ is a G -cover, the **ramification locus** of π is the locus $R := R_\pi \subset X$ of points with non-trivial stabilizer; the **(reduced) branch locus** of π is $\mathcal{B}_\pi := \pi(R) \subset Y$.

In the case where X, Y are smooth, R is the reduced divisor defined by the Jacobian determinant of π , and also \mathcal{B}_π is a divisor (purity of the branch locus).

In the following, if there is no danger of confusion, we will denote \mathcal{B}_π by \mathcal{B} .

Let $\pi: X \rightarrow Y$ be a G -cover. Since $\pi: X \rightarrow Y$ is finite, $\pi_*\mathcal{O}_X$ is a coherent sheaf of \mathcal{O}_Y -algebras. The \mathcal{O}_Y -algebra structure on $\pi_*\mathcal{O}_X$ corresponds to a morphism $m: \pi_*\mathcal{O}_X \otimes_{\mathcal{O}_Y} \pi_*\mathcal{O}_X \rightarrow \pi_*\mathcal{O}_X$ of \mathcal{O}_Y -modules that gives a commutative and associative product on $\pi_*\mathcal{O}_X$. These data determine X as $\text{Spec } \pi_*\mathcal{O}_X$. Furthermore, the G -action on X corresponds to a G -action on $\pi_*\mathcal{O}_X$ which is the identity on \mathcal{O}_Y . The two structures on $\pi_*\mathcal{O}_X$, of an algebra over \mathcal{O}_Y and of a G -module, impose restrictions that sometimes allow one to determine certain “building data” for the G -cover π . One of our aims here is the determination of such building data when $G = D_n$ (Theorem 5.6).

We will restrict ourselves to the case where π is flat, or equivalently $\pi_*\mathcal{O}_X$ is locally free. We recall in the next proposition a useful characterization of flat G -covers.

Proposition 2.2. Let $\pi: X \rightarrow Y$ be a G -cover, with X irreducible and Y smooth. Then π is flat $\Leftrightarrow X$ is Cohen-Macaulay.

Proof. (\Leftarrow) This is a direct application of the Corollary to Theorem 23.1 in [Mat89].

(\Rightarrow) Let $x \in X$ and $y = \pi(x)$. Since π is flat, for any regular sequence $a_1, \dots, a_r \in m_y$ for $\mathcal{O}_{Y,y}$, $\pi^*(a_1), \dots, \pi^*(a_r)$ is a regular sequence for $\mathcal{O}_{X,x}$. Hence, since Y is smooth and π is finite, X is Cohen-Macaulay. \square

In the rest of this section we assume that X and Y are smooth. The aim here is to relate the basic sheaves of X , namely $\Omega_X^i, \Theta_X, \dots$, with the corresponding ones of Y . Concretely, for example for the Ω_X^i 's, there is a chain of inclusions as follows:

$$\Omega_Y^i \otimes \pi_*\mathcal{O}_X \hookrightarrow \pi_*\Omega_X^i \hookrightarrow \Omega_Y^i(\log \mathcal{B})^{**} \otimes \pi_*\mathcal{O}_X,$$

where $\Omega_Y^i(\log \mathcal{B})$ is the sheaf of i -forms with logarithmic poles along \mathcal{B} and $\Omega_Y^i(\log \mathcal{B})^{**}$ is its double dual. These three sheaves coincide on $Y \setminus \mathcal{B}$ and it is possible to describe explicitly $\pi_*\Omega_X^i$ inside $\Omega_Y^i(\log \mathcal{B})^{**} \otimes$

$\pi_*\mathcal{O}_X$ at the generic points \mathcal{B}_i of \mathcal{B} . Then, by Hartogs' theorem, one obtains a description of $\pi_*\Omega_X^1$. We work out in detail the steps of this procedure for Ω_X^1 and Θ_X .

Let $\mathbb{C}(X)$ and $\mathbb{C}(Y)$ be the fields of rational functions on X and Y , respectively. The space of differentials of $\mathbb{C}(X)$, $\Omega_{\mathbb{C}(X)}^1$, is a vector space of dimension

$$d = \dim_{\mathbb{C}(X)} \Omega_{\mathbb{C}(X)}^1 = \dim X$$

and a basis is given by the differentials of a transcendence basis of the field extension $\mathbb{C} \subset \mathbb{C}(X)$. The same holds true for $\Omega_{\mathbb{C}(Y)}^1$ and, since $\dim X = \dim Y$, the pull-back morphism $\pi^*: \mathbb{C}(Y) \rightarrow \mathbb{C}(X)$ induces an isomorphism of $\mathbb{C}(X)$ -vector spaces:

$$(1) \quad \Omega_{\mathbb{C}(Y)}^1 \otimes_{\mathbb{C}(Y)} \mathbb{C}(X) \rightarrow \Omega_{\mathbb{C}(X)}^1.$$

In other words, the rational differentials of X can be written as linear combinations

$$\sum_{i=1}^d f_i dy_i,$$

where $f_1, \dots, f_d \in \mathbb{C}(X)$, $y_1, \dots, y_d \in \mathbb{C}(Y)$ form a transcendence basis for the extension $\mathbb{C} \subset \mathbb{C}(Y)$. Since a regular differential 1-form on X is a rational differential 1-form which is regular at each point, the isomorphism (1) induces an inclusion

$$(2) \quad \varphi: \pi_*\Omega_X^1 \hookrightarrow \Omega_{\mathbb{C}(Y)}^1 \otimes_{\mathbb{C}(Y)} \mathbb{C}(X).$$

Since X and Y are smooth, by the theorem of purity of the branch locus [Zar58] the ramification locus R and the branch locus \mathcal{B} of π are divisors in X and Y respectively. Let $\mathcal{B} = \sum_{i=1}^r \mathcal{B}_i$, where $\mathcal{B}_i \subset Y$ are prime divisors. We recall, following [CHKS06], the following definition:

$$\Omega_Y^1(\log \mathcal{B}) := \text{Im} \left(\Omega_Y^1 \oplus \mathcal{O}_Y^{\oplus r} \rightarrow \Omega_Y^1(\mathcal{B}) \right),$$

where $\Omega_Y^1 \hookrightarrow \Omega_Y^1(\mathcal{B})$ is the natural inclusion, $\mathcal{O}_Y^{\oplus r} \rightarrow \Omega_Y^1(\mathcal{B})$ is given by

$$e_i \mapsto \frac{db_i}{b_i}, \quad i = 1, \dots, r,$$

where, for any i , e_i is the i -th element of the standard \mathcal{O}_Y -basis of $\mathcal{O}_Y^{\oplus r}$, and b_i is a global section of $\mathcal{O}_Y(\mathcal{B}_i)$ with $\mathcal{B}_i = \{b_i = 0\}$.

Let $\Omega_Y^1(\log \mathcal{B})^{**}$ be the double dual of $\Omega_Y^1(\log \mathcal{B})$. It coincides with the sheaf of germs of logarithmic 1-forms with poles along \mathcal{B} defined in [Sai80]. Indeed both sheaves are reflexive and they coincide on the complement $Y \setminus \text{Sing}(\mathcal{B})$ of the singular locus of \mathcal{B} .

Proposition 2.3. *Let $\pi: X \rightarrow Y$ be a G -cover with X and Y smooth. Then there are G -equivariant inclusions of sheaves of \mathcal{O}_Y -modules as follows:*

$$\Omega_Y^1 \otimes \pi_*\mathcal{O}_X \hookrightarrow \pi_*\Omega_X^1 \hookrightarrow \Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_*\mathcal{O}_X,$$

where the morphisms are isomorphisms on the complement $Y \setminus \mathcal{B}$ of the branch divisor.

Moreover, let $\xi_i \in \pi_* \mathcal{O}_X$ be the generator of the ideal of the reduced divisor $R_i := \pi^{-1}(\mathcal{B}_i)$ and a maximal local root of b_i ($b_i = \xi_i^{m_i}$), for any $i = 1, \dots, r$. Then $\pi_* \Omega_X^1$ is characterized as the subsheaf of $\Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_* \mathcal{O}_X$ such that at the smooth points of $\mathcal{B}_i = \{b_i = 0\}$ it coincides with the subsheaf of \mathcal{O}_Y -modules generated by $\Omega_Y^1 \otimes \pi_* \mathcal{O}_X$ and by the elements $\xi_i^k d \log(b_i)$, $k = 1, \dots, m_i - 1$.

Proof. The first arrow to the left is the push-forward under π_* of the natural inclusion

$$(T\pi)^*: \pi^* \Omega_Y^1 \hookrightarrow \Omega_X^1,$$

where $T\pi$ is the tangent map of π .

Now consider the morphism φ in (2). We first show that $(\text{Im}\varphi)_q \subset (\Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_* \mathcal{O}_X)_q$ for any $q \in Y' := Y \setminus \text{Sing}(\mathcal{B})$. On the complement of \mathcal{B} , $Y \setminus \mathcal{B}$, this follows from the projection formula, since π is not ramified on $\pi^{-1}(Y \setminus \mathcal{B})$ and hence

$$\pi_* \Omega_X^1 \cong \Omega_Y^1 \otimes \pi_* \mathcal{O}_X \cong \Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_* \mathcal{O}_X \quad \text{on } Y \setminus \mathcal{B}.$$

Let now $q \in \mathcal{B}' := \mathcal{B} \setminus \text{Sing}(\mathcal{B})$ and $p \in \pi^{-1}(q)$. Choose local coordinates y_1, \dots, y_d at q and x_1, \dots, x_d at p , such that π has the following local expression:

$$y_1 = x_1^m, y_2 = x_2, \dots, y_d = x_d.$$

Hence $\mathcal{O}_{X,p} = \mathcal{O}_{Y,q}[x_1]/(x_1^m - y_1)$, and the stalk $(\Omega_Y^1)_q$ is generated over $\mathcal{O}_{Y,q}$ by

$$d y_1, d y_2, \dots, d y_d,$$

while the stalk $(\Omega_X^1)_p$ is generated over $\mathcal{O}_{X,p}$ by

$$d x_1, d x_2, \dots, d x_d,$$

with the obvious relations

$$m \cdot d \log(x_1) = d \log(y_1), \quad \text{equivalently} \quad d x_1 = \frac{x_1}{m y_1} d y_1,$$

$$d x_i = d y_i, \quad i \geq 2.$$

From this it follows that $(\pi_* \Omega_X^1)_q$ is generated as $\mathcal{O}_{Y,q}$ -module by the elements of the G -orbit of

$$\begin{cases} x_1^i d y_j & 0 \leq i \leq m-1, 2 \leq j \leq d, \\ x_1^k \frac{d y_1}{y_1}, & 1 \leq k \leq m-1 \\ d y_1 \end{cases}$$

which implies the claim at the points $q \in \mathcal{B}'$, since $y_1 = 0$ is a defining equation for \mathcal{B} near q . Moreover, we see that $(\pi_* \Omega_X^1)_q$ is generated, as an \mathcal{O}_Y -module, by $(\Omega_Y^1)_q \otimes (\pi_* \mathcal{O}_X)_q$ and by the elements $x_1^k \frac{d y_1}{y_1}$, $1 \leq k \leq m-1$.

Notice that this argument also shows that the inclusion $(\pi_* \Omega_X^1)_q \subset (\Omega_{Y,q}(\log \mathcal{B})^{**} \otimes (\pi_* \mathcal{O}_X))_q$ is G -equivariant, if $q \in Y \setminus \text{Sing}(\mathcal{B})$.

To conclude the proof, observe that $\pi_*\Omega_X^1$ is locally free and that $\Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_*\mathcal{O}_X$ is reflexive, so that any section of $\pi_*\Omega_X^1$ on $Y \setminus \text{Sing}(\mathcal{B})$ has a unique extension in $\Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_*\mathcal{O}_X$. \square

Let now $\Theta_X := \mathcal{H}om_{\mathcal{O}_X}(\Omega_X^1, \mathcal{O}_X)$ and $\Theta_Y := \mathcal{H}om_{\mathcal{O}_Y}(\Omega_Y^1, \mathcal{O}_Y)$ be the tangent sheaves of X and Y , respectively. In the following proposition we relate the sheaf $\pi_*\Theta_X$ with Θ_Y .

Define as in [Sai80] or [Cat88, Def. 9.15] the sheaf $\Theta_Y(-\log \mathcal{B})$ of logarithmic vector fields on Y with respect to $\mathcal{B} = \{b = 0\}$ as

$$\Theta_Y(-\log \mathcal{B}) := \{v | v \cdot \log(b) \in \mathcal{O}_Y\} = \{v | v(b) \in b\mathcal{O}_Y\} = (\Omega_Y^1(\log \mathcal{B}))^*.$$

Observe that the quotient sheaf $\Theta_Y/\Theta_Y(-\log \mathcal{B})$ equals the equisingular normal sheaf $N'_{\mathcal{B}|Y}$ of \mathcal{B} in Y , which coincides with the normal sheaf at the points where \mathcal{B} is smooth ([Cat88], rem. 9.16).

Let us recall that the usual pairing $\Theta_Y \times \Omega_Y^1 \rightarrow \mathcal{O}_Y$ extends to a perfect pairing

$$\Theta_Y(-\log \mathcal{B}) \times \Omega_Y^1(\log \mathcal{B})^{**} \rightarrow \mathcal{O}_Y.$$

Proposition 2.4. *Let $\pi: X \rightarrow Y$ be a G -cover with X and Y smooth. Let $\mathcal{B} \subset Y$ be the branch divisor of π . Then the tangent map of π identifies $\pi_*\Theta_X$ with a subsheaf of $\Theta_Y \otimes \pi_*\mathcal{O}_X$. Under this identification we have the following inclusions of sheaves*

$$\Theta_Y(-\log \mathcal{B}) \otimes \pi_*\mathcal{O}_X \subset \pi_*\Theta_X \subset \Theta_Y \otimes \pi_*\mathcal{O}_X$$

which are compatible with the G -actions. Moreover the three sheaves coincide on $Y \setminus \mathcal{B}$ and $\pi_*\Theta_X$ is characterized as the subsheaf of $\Theta_Y \otimes \pi_*\mathcal{O}_X$ sending $\pi_*\Omega_X^1$ to $\pi_*\mathcal{O}_X$.

More concretely, it is the subsheaf such that at the smooth points of $\mathcal{B}_i = \{b_i = 0\}$ coincides with the subsheaf of \mathcal{O}_Y -modules generated by $\Theta_Y(-\log \mathcal{B}) \otimes \pi_*\mathcal{O}_X$ and by $\xi_i^{-1}b_i \frac{\partial}{\partial b_i}$, where $\xi_i \in \pi_*\mathcal{O}_X$ is the generator of the ideal of the reduced divisor $R_i := \pi^{-1}(\mathcal{B}_i)$ (and a maximal local root of b_i , $b_i = \xi_i^{m_i}$), while $b_i \frac{\partial}{\partial b_i}$ is a local generator of $N'_{\mathcal{B}|Y}$.

Proof. The tangent morphism $T\pi: \Theta_X \rightarrow \pi^*\Theta_Y$ gives a G -equivariant morphism of sheaves

$$(3) \quad \pi_*\Theta_X \rightarrow \Theta_Y \otimes \pi_*\mathcal{O}_X.$$

We first prove that (3) is injective and that its image contains $\Theta_Y(-\log \mathcal{B}) \otimes \pi_*\mathcal{O}_X$. By Hartogs' theorem and the definition of $\Theta_Y(-\log \mathcal{B})$ it suffices to prove this on the complement of $\text{Sing}(\mathcal{B})$.

On $Y \setminus \mathcal{B}$, π is not ramified, hence (3) is an isomorphism. Let now $q \in \mathcal{B} \setminus \text{Sing}(\mathcal{B})$ and let $p \in \pi^{-1}(q)$. Choose as before coordinates y_1, \dots, y_d at q and x_1, \dots, x_d at p , such that π has the following local expression:

$$y_1 = x_1^m, y_2 = x_2, \dots, y_d = x_d.$$

Then $(\pi_*\Theta_X)_q$ is generated by the G -orbit of

$$x_1^k \left(\frac{\partial}{\partial x_1} + mx_1^{m-1} \frac{\partial}{\partial y_1} \right), x_1^k \frac{\partial}{\partial y_2}, \dots, x_1^k \frac{\partial}{\partial y_d}, 0 \leq k \leq m-1,$$

as $\mathcal{O}_{Y,q}$ -module. The image of these generators under (3) is the G -orbit of

$$m \frac{\partial}{\partial y_1} \otimes x_1^{m-1+k}, \frac{\partial}{\partial y_2} \otimes x_1^k, \dots, \frac{\partial}{\partial y_d} \otimes x_1^k, 0 \leq k \leq m-1.$$

From this it follows that (3) is injective and that its image contains the sheaf $\Theta_Y(-\log \mathcal{B}) \otimes \pi_*\mathcal{O}_X$, which is generated by the G -orbit of

$$y_1 \frac{\partial}{\partial y_1} \otimes x_1^k = \frac{\partial}{\partial y_1} \otimes x_1^{m+k}, \frac{\partial}{\partial y_2} \otimes x_1^k, \dots, \frac{\partial}{\partial y_d} \otimes x_1^k, \quad 0 \leq k \leq m-1,$$

as $\mathcal{O}_{Y,q}$ -module. The only missing term to get from $\Theta_Y(-\log \mathcal{B}) \otimes \pi_*\mathcal{O}_X$ the full direct image is then $y_1 \frac{\partial}{\partial y_1} \otimes x_1^{-1}$. We observe that $y_1 \frac{\partial}{\partial y_1} \otimes x_1^{-1} = \frac{1}{x_1} y_1 \frac{\partial}{\partial y_1}$ and the vector field $y_1 \frac{\partial}{\partial y_1}$ is the local generator of $N'_{\mathcal{B}|Y} = \Theta_Y / \Theta_Y(-\log \mathcal{B})$. \square

Remark 2.5. Notice that the previous results are valid for more general Galois covers of complex manifolds.

2.1. The non-Galois case. We extend propositions 2.3 and 2.4 to branched covers $\pi: X \rightarrow Y$ which are not necessarily Galois. Notice that also in this case we have an inclusion of the sheaf $\pi_*\Omega_X^1$ in the constant sheaf $\Omega_{\mathbb{C}(Y)}^1 \otimes_{\mathbb{C}(Y)} \mathbb{C}(X)$ as in (2).

Proposition 2.6. *Let $\pi: X \rightarrow Y$ be a branched cover with X and Y smooth. Let $\mathcal{B} \subset Y$ be the branch divisor of π . Then we have the following inclusions of \mathcal{O}_Y -modules:*

$$\Omega_Y^1 \otimes \pi_*\mathcal{O}_X \hookrightarrow \pi_*\Omega_X^1 \hookrightarrow \Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_*\mathcal{O}_X,$$

where the morphism on the left is induced by the pull-back morphism $(T\pi)^*: \pi^*\Omega_Y^1 \rightarrow \Omega_X^1$ and the one on the right is induced by (2).

Proof. Since $\pi: X \setminus R \rightarrow Y \setminus \mathcal{B}$ is non-ramified, $(T\pi)^*: \pi^*\Omega_Y^1 \rightarrow \Omega_X^1$ is injective. Applying the push-forward functor π_* we obtain an injective homomorphism $\Omega_Y^1 \otimes \pi_*\mathcal{O}_X \hookrightarrow \pi_*\Omega_X^1$.

We now prove that the morphism (2) has image in $\Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_*\mathcal{O}_X$. Notice that by the same argument as in the proof of Proposition 2.3, it suffices to prove this on the complement of the singular locus of \mathcal{B} . So we assume that \mathcal{B} is smooth. The assertion is true on $Y \setminus \mathcal{B}$, because here π is unramified. Let now $q \in \mathcal{B}$ and let $V \subset Y$ be an open neighbourhood (in the complex topology) of q , such that:

$$\pi^{-1}(V) = U_1 \sqcup \dots \sqcup U_s,$$

with $U_1, \dots, U_s \subset X$ disjoint subsets; for any $i = 1, \dots, s$ there are coordinates $(x_{i,1}, \dots, x_{i,d})$ on U_i and (y_1, \dots, y_d) on V , such that the

restriction of π on U_i has the form

$$(x_{i,1}, \dots, x_{i,d}) \mapsto (y_1 = x_{i,1}^{e_i}, y_2 = x_{i,2}, \dots, y_d = x_{i,d}),$$

e_1, \dots, e_s are integers ≥ 1 (see e.g. [Namba87]). Then $(\pi_* \Omega_X^1)|_V$ is generated as $\mathcal{O}_Y(V)$ -module by:

$$\begin{cases} x_{i,1}^{k_i} dx_{i,1} = \frac{x_{i,1}^{k_i+1}}{e_i} \frac{dy_1}{y_1}, & k_i = 0, \dots, e_i - 1, i = 1, \dots, s \\ x_{i,1}^{k_i} dx_{i,j} = x_{i,1}^{k_i} dy_j, & j = 2, \dots, d, i = 1, \dots, s, k_i = 0, \dots, e_i - 1. \end{cases}$$

Since $y_1 = 0$ is a local equation for \mathcal{B} , it follows that the image of $\pi_* \Omega_X^1$ under (2) is contained in $\Omega_Y^1(\log \mathcal{B})^{**} \otimes \pi_* \mathcal{O}_X$. \square

Analogously, for the tangent sheaves we have the following

Proposition 2.7. *Let $\pi: X \rightarrow Y$ be a branched cover, with X and Y smooth. Let $\mathcal{B} \subset Y$ be the branch locus. Then the tangent map of π induces an injective morphism of sheaves*

$$\pi_* \Theta_X \hookrightarrow \Theta_Y \otimes \pi_* \mathcal{O}_X$$

whose image contains the sheaf $\Theta_Y(-\log \mathcal{B}) \otimes \pi_* \mathcal{O}_X$.

Remark 2.8. We can obtain more precise results by taking the Galois closure $p: W \rightarrow Y$, which is a normal variety, and writing $X = W/H$ where H is a suitable subgroup of the Galois group G . The only problem is that W may not be smooth, and we denote then by W' its smooth locus. Then we can write $\pi_* \Omega_X^1$ as essentially the submodule of H -invariants inside $p_* \Omega_{W'}^1$: for instance we have

$$\pi_* \Omega_X^1 = ((p_* \Omega_{W'}^1)^H)^{**},$$

since the regular 1-forms on $X' = W'/H$ are just the H -invariant 1-forms on W' .

3. LINE BUNDLES AND DIVISORIAL SHEAVES ON FLAT DOUBLE COVERS

In this section we consider the following general situation. We have a flat finite double cover, where Y is smooth:

$$q: Z \rightarrow Y, \quad Z = \text{Spec}(q_* \mathcal{O}_Z),$$

$$\mathcal{R} := q_* \mathcal{O}_Z = (\mathcal{O}_Y \oplus z \mathcal{O}_Y(-L)/(z^2 - F)), \quad F \in H^0(\mathcal{O}_Y(2L)).$$

Our goal is to have a description of divisorial sheaves, that is, rank 1 reflexive sheaves \mathcal{L} on the Gorenstein variety Z in terms of their direct image $q_*(\mathcal{L}) =: \mathcal{N}$.

By flatness of \mathcal{L} over Y , which we assume throughout this section, \mathcal{N} is a rank two vector bundle on Y (a locally free sheaf of rank 2), and its \mathcal{R} -module structure is fully encoded in an endomorphism

$$N: \mathcal{N}(-L) \rightarrow \mathcal{N}$$

such that

$$N^2 = F \cdot \text{Id}_{\mathcal{N}} : \mathcal{N}(-2L) \rightarrow \mathcal{N}.$$

In particular, observe that $\text{Tr}(N) = 0$, $\det(N) = -F$, so that, on any open set where \mathcal{N} and the divisor L are trivialized, N is given by a matrix of the form

$$\begin{pmatrix} a & b \\ c & -a \end{pmatrix}, \quad a^2 + bc = F.$$

Conversely, any such pair (\mathcal{N}, N) as above determines a divisorial sheaf \mathcal{L} on Z .

Lemma 3.1. *Given a pair (\mathcal{N}, N) where $N : \mathcal{N}(-L) \rightarrow \mathcal{N}$ satisfies $N^2 = F \cdot \text{Id}_{\mathcal{N}} : \mathcal{N}(-2L) \rightarrow \mathcal{N}$, it determines a saturated rank 1 torsion free \mathcal{R} -module which is locally free exactly in the points of Y where the endomorphism N does not vanish. In particular, a divisorial sheaf \mathcal{L} on Z .*

Proof. View locally \mathcal{N} as \mathcal{O}_Y^2 : then a local section (x, y) is a local \mathcal{R} generator at a point P if and only if (x, y) and $N(x, y)$ are a local basis. This means, in terms of the matrix N , that (x, y) and $(ax + by, cx - ay)$ are linearly independent; equivalently, the determinant

$$q(x, y) = cx^2 - 2axy - by^2 \neq 0.$$

Hence \mathcal{N} is an invertible \mathcal{R} -module if and only if the quadratic form obtained by evaluating a, b, c at P is not identically zero, which amounts to the condition that a, b, c do not vanish simultaneously. \square

Remark 3.2. Of course, the points where N vanishes yield singular points of the branch locus $\mathcal{B} = \{F = 0\}$, and singular points of Z .

In the case where Z is normal, then \mathcal{L} is determined by its restriction on the smooth locus Z^0 of Z since $\mathcal{L} = i_*(\mathcal{L}|_{Z^0})$ (we are denoting by $i : Z^0 \rightarrow Z$ the inclusion map), hence we are just dealing with the Picard group $\text{Pic}(Z^0)$. We want now to spell out in detail the group structure of $\text{Pic}(Z^0)$ in terms of the direct image rank 2 vector bundles.

Proposition 3.3. *Let Z be normal, let $\mathcal{L}_1^0, \mathcal{L}_2^0$ be invertible sheaves on Z^0 , and let $\mathcal{N}_j = (q \circ i)_*(\mathcal{L}_j^0)$, $j = 1, 2$. Then the tensor product operation in $\text{Pic}(Z^0)$, associating to $\mathcal{L}_1^0, \mathcal{L}_2^0$ their tensor product $\mathcal{L}_1^0 \otimes_{\mathcal{O}_{Z^0}} \mathcal{L}_2^0$ gives the following pair:*

1) the vector bundle $(q \circ i)_*(\mathcal{L}_1^0 \otimes_{\mathcal{O}_{Z^0}} \mathcal{L}_2^0)$ equals the \mathcal{O}_Y -double dual of $\mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2$, which is the cokernel of the following exact sequence:

$$\psi : (\mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2)(-L) \rightarrow \mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2 \rightarrow \mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2 \rightarrow 0,$$

where

$$\psi = N_1 \otimes_{\mathcal{O}_Y} \text{Id}_{\mathcal{N}_2} - \text{Id}_{\mathcal{N}_1} \otimes_{\mathcal{O}_Y} N_2.$$

2) To the inverse invertible sheaf \mathcal{L}^{0-1} corresponds the \mathcal{R} -module associated to the pair $(\mathcal{N}^*(-L), {}^t N(-2L))$, where $\mathcal{N}^* := \mathcal{H}om(\mathcal{N}, \mathcal{O}_Y)$.

3) Two invertible sheaves on Z^0 , $\mathcal{L}_1^0, \mathcal{L}_2^0$, are isomorphic if and only if there is an isomorphism $\Psi : \mathcal{N}_1 \rightarrow \mathcal{N}_2$ such that $\Psi \circ N_1 = N_2 \circ \Psi(-L)$.

4) Effective Weil divisors D on Z correspond to points $[\delta]$ of some projective space $\mathbb{P}(H^0(\mathcal{N}))$, for some isomorphism class of a pair (\mathcal{N}, N) as above. The sum $D_1 + D_2$ corresponds to the image of $\delta_1 \otimes_{\mathcal{R}} \delta_2$ in the double dual $(\mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2)^{**}$.

Proof. 1) First of all we have that $(q \circ i)_*(\mathcal{L}_1^0 \otimes_{\mathcal{O}_{Z^0}} \mathcal{L}_2^0)$ is the saturation of the sheaf equal to $\mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2$ on the open set $Y^0 = Y \setminus \text{Sing}(\mathcal{B})$ (letting $j : Y^0 \rightarrow Y$ be the inclusion, the saturation of \mathcal{F} is here for us $j_*(\mathcal{F}|_{Y^0})$).

Moreover, by definition, $\mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2$ is the quotient of $\mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2$ by the submodule generated by the elements $(z \cdot n_1) \otimes_{\mathcal{O}_Y} n_2 - n_1 \otimes_{\mathcal{O}_Y} (z \cdot n_2)$, i.e., the submodule image of

$$N_1 \otimes_{\mathcal{O}_Y} \text{Id}_{\mathcal{N}_2} - \text{Id}_{\mathcal{N}_1} \otimes_{\mathcal{O}_Y} N_2 : \mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2(-L) \rightarrow \mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2.$$

Since the above is an antisymmetric map of vector bundles, its rank at one point is either 2, or 0: but the latter happens, as it is easy to verify, exactly when both N_1 and N_2 vanish, i.e., in a locus contained in the singular locus of the branch locus $\mathcal{B} = \{F = 0\}$, which has codimension 2 since we assume Z to be normal.

Hence we have that $\mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2$ is a rank 2 bundle at $y \in Y$ if either N_1 or N_2 do not vanish at y ; in the contrary case, the sheaf $\mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2$ has rank 4 at the point y , and one needs to take $(\mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2)^{**}$, where we set $\mathcal{F}^* := \mathcal{H}om_{\mathcal{O}_Y}(\mathcal{F}, \mathcal{O}_Y)$.

We observe moreover that we have an infinite complex

$$\dots \rightarrow (\mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2)(-2L) \rightarrow (\mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2)(-L) \rightarrow \mathcal{N}_1 \otimes_{\mathcal{O}_Y} \mathcal{N}_2 \rightarrow \mathcal{N}_1 \otimes_{\mathcal{R}} \mathcal{N}_2 \rightarrow 0,$$

where the maps are given by

$$\psi' = N_1 \otimes_{\mathcal{O}_Y} \text{Id}_{\mathcal{N}_2} + \text{Id}_{\mathcal{N}_1} \otimes_{\mathcal{O}_Y} N_2,$$

respectively by the above ψ . The complex is exact on the right and also at the points where either N_1 or N_2 are not vanishing.

2) Given the module $\mathcal{H}om_{\mathcal{R}}(\mathcal{N}, \mathcal{R})$, its elements consist of elements $\phi = \phi_1 + z\phi_2$, where $\phi_1 \in \mathcal{N}^*$, and $\phi_2 : \mathcal{N} \rightarrow \mathcal{O}_Y(-L)$. \mathcal{R} -linearity is equivalent to $\phi(Nn) = z\phi(n)$, hence to

$$\phi_2(Nn) = \phi_1(n), \phi_1(Nn) = F\phi_2(n).$$

The first equation implies the second, hence ϕ_2 determines ϕ ; moreover, given any $\phi_2 : \mathcal{N} \rightarrow \mathcal{O}_Y(-L)$, and defining $\phi_1(n) := \phi_2(Nn)$, we get an \mathcal{R} -linear homomorphism by the above formulae.

Clearly, multiplication by z acts on ϕ by sending $\phi = \phi_1 + z\phi_2 \mapsto F\phi_2 + z\phi_1$, hence it sends $\phi_2 : \mathcal{N} \rightarrow \mathcal{O}_Y(-L)$ to $\phi_2 \circ N$, and it is given by ${}^t N(-2L) : \mathcal{N}^*(-L)(-L) \rightarrow \mathcal{N}^*(-L)$.

3) Is easy to show, applying $(q \circ i)_*$ to the given isomorphism.

4) A section of \mathcal{L} determines, again by applying $(q \circ i)_*$, a section $\delta \in H^0(\mathcal{N})$, which represents the image of $1 \in \mathcal{O}_Y$ and which in turn determines the image of an element $\alpha + z\beta \in \mathcal{R}$: since we must have $\alpha + z\beta \mapsto \delta(\alpha) + N\delta(\beta)$.

□

4. LINE BUNDLES ON HYPERELLIPTIC CURVES

We apply the theory developed in the previous section to discuss the particular, but very interesting case, of line bundles on hyperelliptic curves. Here $Y = \mathbb{P}^1$, and $Z = C$ is a hyperelliptic curve of genus $g \geq 1$ defined by the equation

$$z^2 = F(x_0, x_1),$$

where F is a homogeneous polynomial of degree $2g+2$ without multiple roots.

The results of this section are also very relevant in order to show the complexity of the equations defining dihedral covers. In fact, a D_n -covering of \mathbb{P}^1 , $X \rightarrow \mathbb{P}^1$, factors through a hyperelliptic curve C . In the case where $X \rightarrow C$ is étale, X is determined by the choice of a line bundle $L \in \text{Pic}^0(C)$ which is an element of n -torsion. This is the reason why we dedicate special attention to the description of n -torsion line bundles on hyperelliptic curves.

For shorthand notation we write $\mathcal{O}(d) := \mathcal{O}_{\mathbb{P}^1}(d)$ and we observe that every locally free sheaf on \mathbb{P}^1 is a direct sum of some invertible sheaves $\mathcal{O}(d_j)$.

Our first goal here is to parametrize the Picard group of C . The first preliminary remark is that it suffices to parametrize the subsets $\text{Pic}^0(C)$, $\text{Pic}^{-1}(C)$, since for any line bundle \mathcal{L} we can choose an integer d such that

$$\deg(\mathcal{L} \otimes_{\mathcal{O}_C} q^*(\mathcal{O}(d))) \in \{0, -1\}.$$

Let us then consider a line bundle \mathcal{L} of degree zero or -1 . In the case $\mathcal{L} = \mathcal{O}_C$ we know that the corresponding pair is the vector bundle

$$\mathcal{O} \oplus z\mathcal{O}(-g-1), N(\alpha + z\beta) = F\beta + z\alpha,$$

and the corresponding matrix is

$$N = \begin{pmatrix} 0 & F \\ 1 & 0 \end{pmatrix}$$

whose determinant yields a trivial factorization $F = 1 \cdot F$.

If instead we assume that \mathcal{L} is nontrivial, i.e. it has no sections, we can write

$$\mathcal{N} := q_*(\mathcal{L}) = \mathcal{O}(-a) \oplus \mathcal{O}(-b), \quad 0 < a \leq b, \quad a+b = g+1-d, \quad d = \deg(\mathcal{L}).$$

The integer a is equal to

$$\min\{m | H^0(\mathcal{L} \otimes q^*\mathcal{O}(m)) \neq 0\},$$

and it determines a Zariski locally closed stratification of the Picard group. We fix a and $d \in \{0, 1\}$ for the time being, and observe that then the vector bundle \mathcal{N} is uniquely determined, whereas the matrix N (determining the \mathcal{R} -structure) has the form

$$N = \begin{pmatrix} P & f \\ q & -P \end{pmatrix}, \quad -\det(N) = (P^2 + qf) = F.$$

Here,

$$\deg(P) = g + 1, \deg(f) = g + 1 - a + b, \deg(q) = g + 1 + a - b.$$

Recall in fact that, in terms of fibre variables (u, v) , we have

$$zu = Pu + qv, zv = fu - Pv.$$

We see the matrix N as providing a factorization

$$F - P^2 = qf.$$

It follows that N is fully determined by the degree $g + 1$ polynomial P and by a partial factorization of the polynomial $F - P^2 = \prod_1^{2g+2} l_i(x_0, x_1)$ (here the l_i are linear forms), as the product fq of two polynomials of respective degrees $\deg(f) = g + 1 - a + b \geq g + 1$, $\deg(q) = g + 1 + a - b \leq g + 1$. Observe that the choice for the polynomial q (hence of f) is only unique up to a constant $\lambda \in \mathbb{C}^*$, once the partial factorization is fixed.

Since for P general the linear forms l_i are distinct, we have in this case exactly $\frac{(2g+2)!}{(g+1-a+b)!(g+1+a-b)!}$ such possible factorizations.

Denote by $\mathcal{V}(a, b)$ the variety of such matrices: $\mathcal{V}(a, b) = \{(P, f, q) | P^2 + qf = F\}$. The preceding discussion shows that the variety $\mathcal{V}(a, b)$ parametrizing such matrices N is a \mathbb{C}^* bundle over a finite covering of degree $\frac{(2g+2)!}{(g+1-a+b)!(g+1+a-b)!}$ of the affine space \mathbb{C}^{g+2} parametrizing our polynomials P . Hence $\mathcal{V}(a, b)$ is an affine variety of dimension $g + 3$. However, in order to get isomorphism classes of line bundles (elements of the Picard group) we must divide by the adjoint action of the group \mathcal{G} of the automorphisms of the vector bundle $\mathcal{N} := q_*(\mathcal{L}) = \mathcal{O}(-a) \oplus \mathcal{O}(-b)$ which have determinant 1.

For $a = b$ we get $\mathcal{G} = SL(2, \mathbb{C})$, whereas for $a < b$ we get a triangular group of matrices

$$B = \begin{pmatrix} \lambda & \beta \\ 0 & \lambda^{-1} \end{pmatrix},$$

where β is any homogeneous polynomial of degree $b - a$.

We also observe that the stabilizer of a matrix N corresponds to an isomorphism of \mathcal{L} , hence to a scalar $\mu \in \mathbb{C}^*$ whose square equals $\det(B) = 1$, hence $\mu = \pm 1$.

We do not investigate here the GIT stability of the orbits, but just observe that the $\mathcal{V}(a, b)$ yields an open set of dimension g in the case where $a = b$, else it gives a stratum of dimension $g + 1 - (b - a)$.

Recalling that $a + b = g + 1 - d$, for $d = 0$ the open set corresponds to: the case $a = b = \frac{g+1}{2}$ for g odd, and the case $a = \frac{g}{2}, b = \frac{g}{2} + 1$ for g even; similarly for $d = -1$.

Next we investigate the explicit description of tensor powers of an invertible sheaf \mathcal{L} on the hyperelliptic curve C , with two motivations: the first one is to try to get useful results towards the description of dihedral coverings of the projective line, the second in order to describe torsion line bundles on hyperelliptic curves.

Proposition 4.1. *There are exact sequences*

$$0 \rightarrow \mathcal{K}_2 := \mathcal{O}(-a - b - (g + 1)) \rightarrow S^2(q_*\mathcal{L}) \rightarrow q_*(\mathcal{L}^2) \rightarrow 0,$$

$$0 \rightarrow \mathcal{K}_3 := \mathcal{K}_2 \otimes q_*\mathcal{L} \rightarrow S^3(q_*\mathcal{L}) \rightarrow q_*(\mathcal{L}^3) \rightarrow 0,$$

$$0 \rightarrow \mathcal{K}_n := \mathcal{K}_2 \otimes S^{n-2}(q_*\mathcal{L}) \rightarrow S^n(q_*\mathcal{L}) \rightarrow q_*(\mathcal{L}^n) \rightarrow 0.$$

Proof. The local sections of $\mathcal{N} = q_*\mathcal{L}$ can be written as pairs (u, v) . Hence $q_*(\mathcal{L}^2)$ is generated by u^2, uv, v^2 , subject to the relation

$$u(fu - Pv) = u(zv) = z(uv) = v(zu) = v(Pu + qv) \Leftrightarrow$$

$$\Leftrightarrow \Xi := fu^2 - 2Puv - qv^2 = 0.$$

That this is the only relation follows since the kernel \mathcal{K}_2 has rank 1 and first Chern class equal to $-a - b - (g + 1)$.

Similarly, $q_*(\mathcal{L}^3)$ is generated by the cubic monomials u^3, u^2v, uv^2, v^3 , and we can simply multiply the relation Ξ by u , respectively v .

In general, we simply observe that there is a natural morphism

$$q_{\mathcal{L}} : C \rightarrow \text{Proj}(q_*\mathcal{L})$$

given by evaluation, and whose image is the relative quadric $\Gamma := \{(u, v) | \Xi(u, v) = 0\}$.

Recall that, since \mathcal{L} is invertible, the polynomials f, P, q cannot vanish simultaneously, in particular $\Gamma := \{\Xi(u, v) = 0\}$ is irreducible. Moreover the branch locus of $\Gamma \rightarrow \mathbb{P}^1$ is $\{(x_0, x_1) | fq + P^2 = F = 0\}$, therefore Γ is isomorphic to C .

Hence C is the hypersurface in $\mathbb{P}' := \text{Proj}(q_*\mathcal{L})$ defined by $\Xi = 0$, where Ξ is a section of $\mathcal{O}_{\mathbb{P}'}(2) \otimes q^*(\mathcal{K}_2)^{-1}$; hence we obtain the general exact sequence via the pushforward of

$$0 \rightarrow \mathcal{O}_{\mathbb{P}'}(n)(-C) = \mathcal{O}_{\mathbb{P}'}(n - 2) \otimes q^*(\mathcal{K}_2) \rightarrow \mathcal{O}_{\mathbb{P}'}(n) \rightarrow \mathcal{O}_C(n) \rightarrow 0.$$

□

Corollary 4.2. *A line bundle \mathcal{L} of n -torsion on the hyperelliptic curve C corresponds to a pair (\mathcal{N}, N) ,*

$$\mathcal{N} = \mathcal{O}(-a) \oplus \mathcal{O}(-b), \quad 0 < a \leq b, \quad a + b = g + 1,$$

$$N = \begin{pmatrix} P & f \\ q & -P \end{pmatrix}, \quad \det(N) = -(P^2 + qf) = -F.$$

where $\deg(P) = g + 1$, $\deg(f) = 2b$, $\deg(q) = 2a$, such that the following linear map

$$H^0(S^n(\mathcal{O}(a) \oplus \mathcal{O}(b))(-2)) \rightarrow H^0(S^{n-2}(\mathcal{O}(a) \oplus \mathcal{O}(b))(2g)),$$

equal to the dual of $\mathcal{K}_2 \otimes S^{n-2}(q_*\mathcal{L}) \rightarrow S^n(q_*\mathcal{L})$ twisted by (-2) , is not surjective.

Proof. Let \mathcal{L} be a degree zero line bundle on C , so that we have $a + b = g + 1$. The condition that \mathcal{L}^n is trivial is clearly equivalent to the condition $H^0(q_*(\mathcal{L}^n)) \neq 0$. In view of the exact cohomology sequence (here $H^0(S^n(q_*\mathcal{L})) = H^0(S^n(\mathcal{N})) = 0$ since $a, b > 0$):

$$\begin{aligned} 0 \rightarrow H^0(\mathcal{K}_n) = 0 \rightarrow H^0(S^n(q_*\mathcal{L})) = 0 \rightarrow H^0(q_*(\mathcal{L}^n)) \rightarrow \\ \rightarrow H^1(\mathcal{K}_n) \rightarrow H^1(S^n(q_*\mathcal{L})) \rightarrow H^1(q_*(\mathcal{L}^n)) \rightarrow 0, \end{aligned}$$

the condition $H^0(q_*(\mathcal{L}^n)) \neq 0$ is equivalent to the non injectivity of $H^1(\mathcal{K}_n) \rightarrow H^1(S^n(q_*\mathcal{L}))$, equivalently, to the non surjectivity of the homomorphism of Serre dual vector spaces:

$$\text{coker}(H^0(S^n(\mathcal{N}^*)(-2)) \rightarrow H^0(S^{n-2}(\mathcal{N}^*)(2g))) \neq 0 \Leftrightarrow$$

$$\Leftrightarrow \text{coker}(H^0(S^n(\mathcal{O}(a) \oplus \mathcal{O}(b))(-2)) \rightarrow H^0(S^{n-2}(\mathcal{O}(a) \oplus \mathcal{O}(b))(2g))) \neq 0.$$

For $n = 2$ this means that, denoting by $\mathcal{A}[m] = H^0(\mathcal{O}(m))$ the space of homogeneous polynomials of degree m , the linear map

$$\mathcal{A}[2a - 2] \oplus \mathcal{A}[a + b - 2] \oplus \mathcal{A}[2b - 2] \rightarrow \mathcal{A}[2a + 2b - 2]$$

given by the matrix $(f, -2P, q)$ is not surjective (equivalently, the linear map given by the matrix (f, P, q) is not surjective).

Writing in non homogeneous coordinates

$$f = \sum_i^{2b} f_i x^i, \quad P = \sum_i^{a+b} P_i x^i, \quad q = \sum_i^{2a} q_i x^i,$$

this means that the matrix

$$A(f, P, q) = \begin{pmatrix} f_0 & f_1 & \cdots & f_{2b-1} & f_{2b} & 0 & 0 & \cdots & 0 \\ 0 & f_0 & f_1 & \cdots & f_{2b-1} & f_{2b} & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & f_0 & f_1 & \cdots & f_{2b-1} & f_{2b} \\ P_0 & P_1 & \cdots & P_{a+b} & 0 & 0 & 0 & \cdots & 0 \\ 0 & P_0 & P_1 & \cdots & P_{a+b} & 0 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & 0 & P_0 & P_1 & \cdots & P_{a+b} \\ q_0 & q_1 & \cdots & q_{2a} & 0 & 0 & 0 & \cdots & 0 \\ 0 & q_0 & q_1 & \cdots & q_{2a} & 0 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & 0 & q_0 & q_1 & \cdots & q_{2a} \end{pmatrix},$$

does not have maximal rank. □

Remark 4.3. The reader may notice the similarity of the matrix $A(f, P, q)$ with the matrix giving the resultant of two homogenous polynomials in two variables. Moreover, notice that $A(f, P, q)$ is a $(3a + 3b - 3) \times (2a + 2b - 1)$ matrix, hence the condition that its rank be at most $2a + 2b - 2$ amounts to a codimension $g = a + b - 1$ condition, which is the expected codimension of the set of n -torsion points in $\text{Pic}^0(C)$.

4.1. Powers of divisorial sheaves on double covers. We would now like to show how the result of proposition 4.1 generalizes to any flat double cover $q : Z \rightarrow Y$. Let \mathcal{L} be a divisorial sheaf on Z , and (\mathcal{N}, N) the associated pair, where $\mathcal{N} = q_*(\mathcal{L})$. There is a natural map $q_{\mathcal{L}} : Z \dashrightarrow \text{Proj}(\mathcal{N})$ given by evaluation, and which is a morphism on the smooth locus of Z .

Set $\mathbb{P}' := \text{Proj}(\mathcal{N})$, and let Γ be the image of $q_{\mathcal{L}}$. Clearly Z is birational to Γ , which is a Gorenstein variety, since it is a divisor in \mathbb{P}' , and $q_{\mathcal{L}}$ is bijective outside of the inverse image of the branch locus \mathcal{B}_q . Moreover q factors through $q_{\mathcal{L}}$ and the natural projection $\pi : \mathbb{P}' \rightarrow Y$.

At the points where \mathcal{L} is invertible, then \mathcal{L} is isomorphic with \mathcal{O}_Z , with an isomorphism compatible with the projection q , hence we conclude that $q_{\mathcal{L}}$ is an embedding on the smooth locus of Z .

At a point P where \mathcal{L} is not invertible, we use the local description of the pair (\mathcal{N}, N) as (\mathcal{O}_Y^2, N) , where N is the matrix sending two local generators x, y to $ax + cy$, respectively to $bx - ay$. Since \mathcal{L} is not invertible, by lemma 3.1 we get that a, b, c vanish at P . We use now the relations

$$xz = ax + cy \Leftrightarrow x(z - a) = cy, \quad yz = bx - ay \Leftrightarrow y(z + a) = bx$$

to obtain

$$(x : y) = (c : z - a) = (z + a : b).$$

These formulae (observe that $(c : z - a) = (z + a : b)$ since $z^2 - a^2 = bc$) show that, at the points P where a, b, c (hence also z) vanish, the rational map $q_{\mathcal{L}}$ blows up the point $q^{-1}(P)$ to the whole fibre \mathbb{P}^1 lying over P . Hence Γ is a small resolution of Z , and we have that the inverse of $q_{\mathcal{L}}$ is obtained blowing down these \mathbb{P}^1 's lying over such points P .

Since we have an isomorphism $Z^0 \cong \Gamma^0$, we see that the line bundle $\mathcal{O}_{\Gamma}(1)$ restricts to \mathcal{L} on Z^0 , hence the sections of \mathcal{L}^n on Z^0 correspond to sections of $\mathcal{O}_{\Gamma}(n)$ on Γ^0 .

Notice that $\text{Pic}(\mathbb{P}^r)$ is generated by $\text{Pic}(Y)$ and by $\mathcal{O}_{\mathbb{P}^r}(1)$, hence there is an invertible sheaf \mathcal{K}_2 on Y such that Γ is the zero set of a section of $\mathcal{O}_{\mathbb{P}^r}(1) \otimes \pi^*(\mathcal{K}_2)^{-1}$.

Consider now the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^r}(n)(-\Gamma) = \mathcal{O}_{\mathbb{P}^r}(n-2) \otimes \pi^*(\mathcal{K}_2) \rightarrow \mathcal{O}_{\mathbb{P}^r}(n) \rightarrow \mathcal{O}_{\Gamma}(n) \rightarrow 0.$$

Taking the direct image, and observing that $\pi_*\mathcal{O}_{\Gamma}(n) = q_*(\mathcal{L}^n)$, we obtain the following

Proposition 4.4. *For each divisorial sheaf \mathcal{L} on Z there is an exact sequence*

$$0 \rightarrow \mathcal{K}_n := \mathcal{K}_2 \otimes S^{n-2}(q_*\mathcal{L}) \rightarrow S^n(q_*\mathcal{L}) \rightarrow q_*(\mathcal{L}^n) \rightarrow 0.$$

5. DIHEDRAL FIELD EXTENSIONS AND GENERALITIES ON DIHEDRAL COVERS

In this section we describe dihedral field extensions $\mathbb{C}(Y) \subset \mathbb{C}(X)$ in the case where Y is factorial and X is normal, thus we obtain a birational classification of D_n -covers. Recall that for any G -cover $\pi: X \rightarrow Y$ of normal varieties the field extension $\mathbb{C}(Y) \subset \mathbb{C}(X)$ is Galois with group G (a G -extension); on the other hand any such field extension determines π as the normalization of Y in $\mathbb{C}(X)$.

In the second part of this section we determine the geometric building data that make the above normalization process explicit. This is important, for instance to calculate invariants of X , to determine the direct images of basic sheaves on X (e.g. $\pi_*\mathcal{O}_X, \pi_*\Omega_X, \pi_*\Theta_X$, etc.) and to provide explicitly families of such covers.

5.1. Dihedral field extensions. Let us fix the following presentation for the dihedral groups:

$$D_n = \langle \sigma, \tau \mid \sigma^n = \tau^2 = (\sigma\tau)^2 = 1 \rangle.$$

The split exact sequence

$$0 \rightarrow \langle \sigma \rangle = \mathbb{Z}/n\mathbb{Z} \rightarrow D_n \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 0$$

gives a factorization of the D_n -cover $\pi: X \rightarrow Y$ as the composition of two intermediate cyclic covers: $\pi = q \circ p$, where $p: X \rightarrow Z$ is a $\mathbb{Z}/n\mathbb{Z}$ -cover, $q: Z \rightarrow Y$ is a $\mathbb{Z}/2\mathbb{Z}$ -cover, $Z := X/\langle \sigma \rangle$. If X is normal and irreducible, then also Z and Y are so, and we have the following chain of field extensions:

$$\mathbb{C}(Y) \subset \mathbb{C}(Z) \subset \mathbb{C}(X),$$

where the field of rational functions on Z is the field $\mathbb{C}(X)^{\langle \sigma \rangle}$ of invariant functions under σ .

Let us recall, following [Cat10], the structure of cyclic extensions $\mathbb{C}(W) \subset \mathbb{C}(V)$, where W, V are normal varieties. Here the Galois group G is cyclic of order m , $G \cong \mathbb{Z}/m\mathbb{Z}$; later we use this description with $m = 2$ and with $m = n$ to study D_n -field extensions. Let $\sigma \in G$ be a generator and let $\zeta \in \mathbb{C}$ be a primitive m -th root of the unity. Then there exists $v \in \mathbb{C}(V)$ and $f \in \mathbb{C}(W)$, such that

$$(4) \quad \begin{cases} \mathbb{C}(V) &= \mathbb{C}(W)(v), & v^m = f, \\ \sigma(v) &= \zeta v. \end{cases}$$

Concretely, v can be chosen to be any non-zero element of the form

$$v = \sum_{i=0}^{m-1} \zeta^{m-i} \sigma^i(\tilde{v}), \quad \tilde{v} \in \mathbb{C}(V).$$

Furthermore, it is possible to choose v and f , such that (4) holds and

$$(5) \quad f = \prod_{i=1}^{m-1} (\delta_i)^i,$$

where $\delta_1, \dots, \delta_{m-1}$ are regular sections of invertible sheaves on $W \setminus \text{Sing}(W)$. To see this, let \hat{v} and \hat{f} satisfying (4), and consider the Weil divisor associated to \hat{f} , $(\hat{f}) = \sum_U v_U(\hat{f})U$, where $U \subset W$ varies among the prime divisors of W and v_U is the valuation of U . On the non-singular locus $W \setminus \text{Sing}(W)$, $U \cap (W \setminus \text{Sing}(W))$ is an effective Cartier divisor, hence $U \cap (W \setminus \text{Sing}(W)) = \{\delta_U = 0\}$, where δ_U is a regular section of an invertible sheaf on $W \setminus \text{Sing}(W)$, hence

$$\hat{f} = \prod_U \delta_U^{v_U(\hat{f})}.$$

Let now $v_U(\hat{f}) = mq_U(\hat{f}) + r_U(\hat{f})$, where $q_U(\hat{f}), r_U(\hat{f}) \in \mathbb{Z}$, $0 \leq r_U(\hat{f}) < m$, and define

$$\delta_i = \prod_{r_U(\hat{f})=i} \delta_U, \quad i = 1, \dots, m-1;$$

then the claim holds true with $v := \hat{v} \prod_U (\delta_U)^{-q_U(\hat{f})}$.

Remark 5.1. If W is factorial, the group of Weil divisors coincides with that of Cartier divisors, hence δ_i corresponds to a Weil divisor D_i which is reduced but not necessarily irreducible. Geometrically D_i is the divisorial part of the branch locus $\mathcal{B} = \sum_{i=1}^{m-1} D_i$ where the local monodromy is σ^i and v is a rational section of a line bundle L on W which satisfies the linear equation

$$(6) \quad mL \equiv \sum_{i=1}^{m-1} iD_i.$$

Conversely, one can construct in a natural way a $\mathbb{Z}/m\mathbb{Z}$ -cover starting from a line bundle L and effective reduced divisors without common components D_1, \dots, D_{m-1} , such that (6) holds ([Par91], [Cat10]).

The following proposition describes dihedral field extensions (see also [Tok94]).

Proposition 5.2. *Let $\mathbb{C}(Y) \subset \mathbb{C}(X)$ be a D_n -extension. Then there exist $a, F \in \mathbb{C}(Y)$ and $x \in \mathbb{C}(X)$, such that*

$$\mathbb{C}(X) = \mathbb{C}(Y)(x), \quad x^{2n} - 2ax^n + F^n = 0,$$

with D_n -action given as follows: $\sigma(x) = \zeta x$, $\tau(x) = F/x$, where $\zeta \in \mathbb{C}$ is a primitive n -th root of 1.

Conversely, given $a, F \in \mathbb{C}(Y)$, such that $x^{2n} - 2ax^n + F^n \in \mathbb{C}(Y)[x]$ is irreducible, then $\frac{\mathbb{C}(Y)[x]}{(x^{2n} - 2ax^n + F^n)}$ is a D_n -field extension of $\mathbb{C}(Y)$ with D_n -action given as before. Hence the normalization of Y in $\frac{\mathbb{C}(Y)[x]}{(x^{2n} - 2ax^n + F^n)}$ is a D_n -covering of Y .

Proof. Consider the quotient $Z := X/\langle\sigma\rangle$ of X by the cyclic subgroup $\langle\sigma\rangle \cong \mathbb{Z}/n\mathbb{Z}$ and let $q: Z \rightarrow Y$ be the induced double cover. From the previous description of cyclic field extensions, we have that

$$\begin{cases} \mathbb{C}(Z) &= \mathbb{C}(Y)(z), \quad z^2 = f \in \mathbb{C}(Y) \\ \bar{\tau}(z) &= -z \end{cases}$$

where $\bar{\tau} \in D_n/\langle\sigma\rangle$ is the class of τ . Since Y is factorial we can assume that f is a regular section of an invertible sheaf on Y .

Now consider the extension $\mathbb{C}(Z) \subset \mathbb{C}(X)$. Let $x \in \mathbb{C}(X)$, $g \in \mathbb{C}(Z)$, such that

$$\begin{cases} \mathbb{C}(X) &= \mathbb{C}(Z)(x), \quad x^n = g, \\ \sigma(x) &= \zeta x, \end{cases}$$

where $\zeta \in \mathbb{C}$ is a primitive n -th root of 1, and notice that $g \in \mathbb{C}(Z)$ can be written as

$$g = a + zb, \quad \text{with } a, b \in \mathbb{C}(Y).$$

Without loss of generality $b = 1$ in the previous formula, otherwise we replace f with $b^2 f$ and z with bz . Moreover, $x\tau(x)$ is invariant under the action of D_n , therefore

$$x\tau(x) = F \in \mathbb{C}(Y), \quad \text{equivalently } \tau(x) = F/x.$$

Finally, $F^n = x^n \tau(x^n) = g\bar{\tau}(g) = (a+z)(a-z) = a^2 - f$, hence

$$f = a^2 - F^n.$$

To conclude, we observe that $z \in \mathbb{C}(Y)(x)$ because $x^n = a + z$, so $\mathbb{C}(X) = \mathbb{C}(Y)(x)$ and by construction it follows that

$$x^{2n} - 2ax^n + F^n = 0.$$

For the last statement, notice that the field extension

$$\mathbb{C}(Y) \subset \frac{\mathbb{C}(Y)[x]}{(x^{2n} - 2ax^n + F^n)}$$

is Galois with group D_n , indeed the conjugates of x , namely $\zeta^i x$ and $\zeta^i F/x$, for $i = 0, \dots, n-1$, belong to $\frac{\mathbb{C}(Y)[x]}{(x^{2n} - 2ax^n + F^n)}$. \square

5.2. Structure of D_n -covers. Let $\pi: X \rightarrow Y$ be a flat D_n -cover with Y smooth. Then $\pi_* \mathcal{O}_X$ is a locally free sheaf of \mathcal{O}_Y -modules, i.e. a vector bundle on Y . Recall that the sheaf $\pi_* \mathcal{O}_X$ carries a natural structure of \mathcal{O}_Y -algebras, which is given by the product of regular functions on X ,

$$(7) \quad m: \pi_* \mathcal{O}_X \otimes_{\mathcal{O}_Y} \pi_* \mathcal{O}_X \rightarrow \pi_* \mathcal{O}_X;$$

the D_n -action on X gives to $\pi_* \mathcal{O}_X$ the structure of a D_n -sheaf (see below). On the other hand, the variety X is completely determined by $\pi_* \mathcal{O}_X$ and m , since $X = \text{Spec}(\pi_* \mathcal{O}_X)$ ([Hart77, II, 5.17]); the D_n -action on X is given by the structure of D_n -sheaf on $\pi_* \mathcal{O}_X$.

Recall that, for any finite group G the *regular representation* of G , $\mathbb{C}[G]$, is the vector space with a basis $\{e_g\}_{g \in G}$ indexed by the elements of G ; for any $h \in G$, an endomorphism of $\mathbb{C}[G]$ is defined by $e_g \mapsto e_{hg}$, for all $g \in G$. In a similar way one defines a sheaf of \mathcal{O}_Y -algebras $\mathcal{O}_Y[G]$, for any variety Y . A sheaf \mathcal{F} of \mathcal{O}_Y -modules is a G -**sheaf**, if it has a structure of sheaf of $\mathcal{O}_Y[G]$ -modules. If moreover \mathcal{F} is a vector bundle, then its fibers carry a linear G -action and so we can see \mathcal{F} as a family of representations of G parametrized by Y .

For any representation $\rho: G \rightarrow \text{GL}(V)$ of G , its *canonical decomposition* (see [Serre77, §2.6]) is the unique decomposition

$$V = V_1 \oplus \dots \oplus V_N$$

defined as follows. Let W_1, \dots, W_N be the different irreducible representations of G . Then each V_i is the direct sum of all the irreducible representations of G in V that are isomorphic to W_i . If χ_1, \dots, χ_N are the characters of W_1, \dots, W_N , and $n_i = \dim W_i$, then

$$(8) \quad p_i = \frac{n_i}{|G|} \sum_{g \in G} \overline{\chi_i(g)} \rho(g) \in \text{End}(V)$$

is the projection of V onto V_i , for any $i = 1, \dots, N$, where $\overline{\chi_i(g)}$ is the complex-conjugate of $\chi_i(g)$.

Let now \mathcal{F} be a locally free G -sheaf on Y with action $\rho: G \rightarrow \text{GL}_{\mathcal{O}_Y}(\mathcal{F})$. Via (8) we define an endomorphism $p_i \in \text{End}_{\mathcal{O}_Y}(\mathcal{F})$, for any $i = 1, \dots, N$. Setting $\mathcal{F}_i := \text{Im}(p_i)$, we have the following decomposition:

$$\mathcal{F} = \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_N.$$

For any $i = 1, \dots, N$, \mathcal{F}_i is the *eigensheaf* of \mathcal{F} corresponding to the (irreducible) representation with character χ_i . Notice that \mathcal{F}_i is a vector sub-bundle of \mathcal{F} , for all i . In particular, when $\mathcal{F} = \pi_* \mathcal{O}_X$, $\pi: X \rightarrow Y$ is a flat G -cover, we have that $\pi_* \mathcal{O}_X$ is a locally free sheaf of $\mathcal{O}_Y[G]$ -modules of rank one, i.e. the fibres of $\pi_* \mathcal{O}_X$ are isomorphic to $\mathbb{C}[G]$ as G -representations. Indeed, the previous procedure gives the decomposition $\pi_* \mathcal{O}_X = (\pi_* \mathcal{O}_X)_1 \oplus \dots \oplus (\pi_* \mathcal{O}_X)_N$, with $(\pi_* \mathcal{O}_X)_i \subset \pi_* \mathcal{O}_X$ a sub-bundle, for any i . By construction, the fibres of $(\pi_* \mathcal{O}_X)_i$ are isomorphic to each other as G -representations. So, it is enough to consider the restriction of $\pi_* \mathcal{O}_X$ on the complement $Y \setminus \mathcal{B}$ of the branch divisor \mathcal{B} , where the assertion follows easily.

In order to describe the sheaves $(\pi_* \mathcal{O}_X)_i$, $i = 1, \dots, N$, when $\pi: X \rightarrow Y$ is a flat D_n -cover, let us briefly recall the representation theory of the dihedral groups. They depend on the parity of n .

n odd. There are two irreducible representations of degree 1 with characters χ_1 and χ_2 ,

$$\begin{array}{c|c|c} & \sigma^k & \sigma^k \tau \\ \hline \chi_1 & 1 & 1 \\ \hline \chi_2 & 1 & -1 \end{array},$$

and $\frac{n-1}{2}$ irreducible representations of degree 2,

$$(9) \quad \rho^\ell(\sigma^k) = \begin{pmatrix} \zeta^{k\ell} & 0 \\ 0 & \zeta^{-k\ell} \end{pmatrix}, \quad \rho^\ell(\sigma^k \tau) = \begin{pmatrix} 0 & \zeta^{k\ell} \\ \zeta^{-k\ell} & 0 \end{pmatrix},$$

where $\zeta \in \mathbb{C}^*$ is a primitive n -th root of 1, $1 \leq \ell \leq \frac{n-1}{2}$, $k = 0, \dots, n-1$.

n even. In this case there are 4 representations of degree 1 with characters χ_1, χ_2, χ_3 and χ_4 ,

	σ^k	$\sigma^k \tau$
χ_1	1	1
χ_2	1	-1
χ_3	$(-1)^k$	$(-1)^k$
χ_4	$(-1)^k$	$(-1)^{k+1}$

and, for any $1 \leq \ell \leq \frac{n}{2} - 1$, the irreducible representation ρ^ℓ defined by (9).

As a consequence we have that

$$\pi_* \mathcal{O}_X = \mathcal{O}_Y \oplus \mathcal{L} \oplus_{\ell=1}^{\frac{n-1}{2}} (\pi_* \mathcal{O}_X)_\ell, \quad \text{if } n \text{ is odd,}$$

and

$$\pi_* \mathcal{O}_X = \mathcal{O}_Y \oplus \mathcal{L} \oplus \mathcal{M} \oplus \mathcal{N} \oplus_{\ell=1}^{\frac{n}{2}-1} (\pi_* \mathcal{O}_X)_\ell, \quad \text{if } n \text{ is even,}$$

where $\mathcal{L}, \mathcal{M}, \mathcal{N}$ are the line bundles corresponding to the 1-dimensional representations with characters respectively χ_2, χ_3, χ_4 .

Notice that the sections of \mathcal{L} are invariant under the action of the subgroup $\langle \sigma \rangle \cong \mathbb{Z}/n\mathbb{Z} \leq D_n$, hence they are regular functions on Z . Moreover we have:

$$(10) \quad q_* \mathcal{O}_Z = \mathcal{O}_Y \oplus \mathcal{L}, \quad \mathcal{L}^{\otimes 2} \cong \mathcal{O}_Y(-\mathcal{B}_q),$$

where $\mathcal{B}_q \subset Y$ is the branch divisor of q .

If n is even, the line bundles \mathcal{M} and \mathcal{N} have a similar interpretation, they arise from the $(\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z})$ -cover $X/\langle \sigma^2, \tau \rangle \rightarrow Y$.

In order to get further information on the rank 4 vector bundles $(\pi_* \mathcal{O}_X)_\ell$, we assume that X is normal and use the factorization $\pi = q \circ p$ and the theory of cyclic covers. Let us denote with Z^0 the smooth locus of Z , $X^0 = p^{-1}(Z^0)$ and $p^0: X^0 \rightarrow Z^0$ be the restriction of p . From the structure theorem for cyclic covers ([Par91], [BaCa08], [Cat10]) it follows that p^0 is determined by divisor classes L_1, \dots, L_{n-1} and reduced effective divisors $D_1^0, \dots, D_{n-1}^0 \subset Z^0$ without common components, such that

$$(11) \quad L_i + L_j \equiv L_{\overline{i+j}} - \sum_{k=1}^{n-1} \varepsilon_{i,j}^k D_k^0,$$

where $\overline{i+j} \in \{0, \dots, n-1\}$, $\overline{i+j} = i+j \pmod{n}$, $L_0 := \mathcal{O}_{Z^0}$.

Let us briefly recall the geometric interpretation of the previous data. For any $i = 0, \dots, n-1$, the line bundle $\mathcal{O}(L_i)$ is the subsheaf of $(p^0)_* \mathcal{O}_{X^0}$ consisting of the regular functions f on X^0 such that $\sigma^* f = \exp\left(\frac{2\pi\sqrt{-1}}{n} i\right) f$. For any $k = 1, \dots, n-1$, the divisor $D_k^0 \subset Z^0$ is the union of the components Δ of the branch divisor \mathcal{B}_{p^0} of p^0 such that, for any component $T \subset (p^0)^{-1}(\Delta)$, the stabilizer of the generic

point of T is the cyclic subgroup of $\langle \sigma \rangle$ generated by σ^k and there is a uniformizing parameter $x \in \mathcal{O}_{X^0, T}$ such that $(\sigma^k)^* x = \exp\left(\frac{2\pi\sqrt{-1}}{|\langle \sigma^k \rangle|}\right) x$, where $|\langle \sigma^k \rangle|$ is the order of σ^k . For every $i, j = 0, \dots, n-1$, $\varepsilon_{i,j}^k$ is defined as follows: let $\nu_i(k), \nu_j(k) \in \{0, \dots, |\langle \sigma^k \rangle| - 1\}$ be such that

$$\begin{aligned} \exp\left(\frac{2\pi\sqrt{-1}}{n} ik\right) &= \exp\left(\frac{2\pi\sqrt{-1}}{|\langle \sigma^k \rangle|}\right)^{\nu_i(k)}, \\ \exp\left(\frac{2\pi\sqrt{-1}}{n} jk\right) &= \exp\left(\frac{2\pi\sqrt{-1}}{|\langle \sigma^k \rangle|}\right)^{\nu_j(k)} \end{aligned}$$

respectively, then

$$(12) \quad \varepsilon_{i,j}^k = \begin{cases} 1 & , \quad \text{if } \nu_i(k) + \nu_j(k) \geq |\langle \sigma^k \rangle| \\ 0 & , \quad \text{otherwise.} \end{cases}$$

Proposition 5.3. *Let Y be a smooth variety and let $\pi: X \rightarrow Y$ be a flat D_n -cover with X normal. Let $p: X \rightarrow Z$ and $q: Z \rightarrow Y$ be the intermediate cyclic covers defined previously. Then the following holds true.*

- (i) $p_* \mathcal{O}_X = \bigoplus_{i=0}^{n-1} \mathcal{F}_i$, where $\mathcal{F}_0 = \mathcal{O}_Z$ and, for any $i = 1, \dots, n-1$, \mathcal{F}_i is a divisorial sheaf on Z . For any $i, j = 1, \dots, n-1$ the product (7) induces an isomorphism as follows:

$$(13) \quad (\mathcal{F}_i \otimes_{\mathcal{O}_Z} \mathcal{F}_j)^{**} \cong \left(\mathcal{F}_{\overline{i+j}} \otimes_{\mathcal{O}_Z} \mathcal{O}_Z \left(- \sum_{k=1}^{n-1} \varepsilon_{i,j}^k D_k \right) \right)^{**},$$

where $\overline{i+j} \in \{0, \dots, n-1\}$, $\overline{i+j} = i+j \pmod{n}$, for any $k = 1, \dots, n-1$, $D_k = \overline{D_k^0}$ is the closure of the divisor D_k^0 in (11), and $()^{**}$ is the double dual in the category of \mathcal{O}_Z -modules.

- (ii) For any $i = 1, \dots, n-1$, $U_i := q_*(\mathcal{F}_i)$ is a vector bundle of rank 2 on Y , in particular \mathcal{F}_i is flat over Y . Furthermore,

$$(\pi_* \mathcal{O}_X)_\ell = U_\ell \oplus U_{n-\ell},$$

where $\ell = 1, \dots, \frac{n-1}{2}$, if n is odd, and $\ell = 1, \dots, \frac{n}{2} - 1$ if n is even; in the case where n is even, $U_{\frac{n}{2}} = \mathcal{M} \oplus \mathcal{N}$.

Proof. Consider the following Cartesian diagram:

$$(14) \quad \begin{array}{ccc} X^0 & \xrightarrow{\iota} & X \\ p^0 \downarrow & & \downarrow p \\ Z^0 & \xrightarrow{j} & Z \end{array}$$

where Z^0 is the smooth locus of Z , $j: Z^0 \rightarrow Z$ is the inclusion, $X^0 = p^{-1}(Z^0)$, $\iota: X^0 \rightarrow X$ is the inclusion, and $p^0 = p|_{X^0}$. Notice that under our hypotheses Z is normal.

Since $\mathcal{O}_X = \iota_* \mathcal{O}_{X^0}$ and $\mathcal{O}_Z = j_* \mathcal{O}_{Z^0}$ (see, e.g. [ReidC3f]),

$$\begin{aligned}
 p_* \mathcal{O}_X &= p_* \iota_* \mathcal{O}_{X^0} = (p \circ \iota)_* \mathcal{O}_{X^0} = \\
 (15) \quad &= j_* p_*^0 \mathcal{O}_{X^0} = j_* (\mathcal{O}_{Z^0} \oplus_{i=1}^{n-1} \mathcal{O}_{Z^0}(L_i)) = \\
 &= \mathcal{O}_Z \oplus_{i=1}^{n-1} j_* \mathcal{O}_{Z^0}(L_i).
 \end{aligned}$$

Let us define $\mathcal{F}_i := j_* \mathcal{O}_{Z^0}(L_i)$, for any $i = 1, \dots, n-1$. The product (7) gives morphisms $m_{ij}: \mathcal{F}_i \otimes \mathcal{F}_j \rightarrow \mathcal{F}_{i+j}$. Since \mathcal{F}_{i+j} is reflexive, m_{ij} is determined by its restriction on the smooth locus Z^0 . On Z^0 , by (11), m_{ij} gives an isomorphism $\mathcal{O}_{Z^0}(L_i + L_j) \cong \mathcal{O}_{Z^0}(L_{i+j} - \sum_{k=1}^{n-1} D_k^0)$. This implies (13).

To prove (ii), apply q_* to (15) and use the following equalities: $\pi = q \circ p$, $q_* \mathcal{O}_Z = \mathcal{O}_Y \oplus \mathcal{L}$. Notice that, $q_* \mathcal{F}_i$ is locally free for any $i = 0, \dots, n-1$, since $\pi_* \mathcal{O}_X$ is locally free and $\pi_* \mathcal{O}_X = \bigoplus_{i=0}^{n-1} q_* \mathcal{F}_i$. The equality $(\pi_* \mathcal{O}_X)_\ell = U_\ell \oplus U_{n-\ell}$ follows directly from the definition of $(\pi_* \mathcal{O}_X)_\ell \subset \pi_* \mathcal{O}_X$ as the eigensheaf corresponding to the representation ρ^ℓ . Finally, in the case where n is even, τ^* acts on $U_{\frac{n}{2}}$; \mathcal{M} is the τ^* -invariant subsheaf, while \mathcal{N} is the τ^* -anti-invariant one. \square

Notice that (13) implies the following relation between \mathcal{F}_1 and the Weil divisors D_1, \dots, D_{n-1} ,

$$(\mathcal{F}_1^{\otimes n})^{**} \cong \mathcal{O}_Z(-\sum_{k=1}^{n-1} kD_k),$$

which is the analogous of Remark 5.1 and equation (6) when the base of the cover is normal.

For later use we observe that the branch divisor $\mathcal{B}_{p^0} = \sum_{k=1}^{n-1} D_k^0$ of $p^0: X^0 \rightarrow Z^0$ is invariant under the involution $\bar{\tau}: Z^0 \rightarrow Z^0$ induced by τ , that is $\bar{\tau}^* \mathcal{B}_{p^0} = \mathcal{B}_{p^0}$, since $\sigma\tau = \tau\sigma^{-1}$. So, there exists an effective Cartier divisor $\Delta_{p^0} \subset Y^0$, such that $(q^0)^* \Delta_{p^0} = \mathcal{B}_{p^0}$, where $q^0 := q|_{Z^0}$. Let us define

$$(16) \quad \Delta_p := \overline{\Delta_{p^0}} \in \text{Div}(Y).$$

Remark 5.4. Notice that, for any $i = 1, \dots, n-1$, $\bar{\tau}(D_i) = D_{n-i}$, so $\bar{\tau}(D_i)$ does not have any common component with D_i for any $i < n/2$, while, if n is even, $\bar{\tau}(D_{\frac{n}{2}}) = D_{\frac{n}{2}}$.

In the remaining of the section we determine building data for D_n -covers of Y . To this aim we first derive some properties of the vector bundles U_1, \dots, U_{n-1} (**1.–3.** below).

1. For any $i = 1, \dots, n-1$, the involution $\tau: X \rightarrow X$ induces an isomorphism $\tau^*: U_i \rightarrow U_{n-i}$, such that $\tau^* \circ \tau^*$ is the identity.

Proof. For any open $V \subset Y$, $U_i(V) := \mathcal{F}_i(q^{-1}(V))$ consists of the regular functions $f \in \mathcal{O}_X(\pi^{-1}(V))$ such that $\sigma^* f = \exp(\frac{2\pi\sqrt{-1}}{n}i)f$. From

the relation $\sigma\tau = \tau\sigma^{-1}$ it follows that $f \mapsto \tau^*f$ gives a morphism $\tau^*: U_i(V) \rightarrow U_{n-i}(V)$ of \mathcal{O}_Y -modules. Since $\tau^2 = 1$, $\tau^* \circ \tau^* = \text{Id}$ and τ^* is an isomorphism. \square

2. For any $i = 1, \dots, n-1$, U_i has a structure of \mathcal{R} -module given by the restriction $m_i: \mathcal{L} \otimes U_i \rightarrow U_i$ of the product (7), where $\mathcal{R} = \mathcal{O}_Y \oplus \mathcal{L} = q_*\mathcal{O}_Z$ (see Section 3.). In particular $m_i^2 = \delta_q \cdot \text{Id}_{U_i}$, where $\delta_q \in H^0(Y, \mathcal{O}_Y(\mathcal{B}_q))$ is such that $\mathcal{B}_q = \{\delta_q = 0\}$ ($\delta_q = F$ of Section 3). Furthermore, the isomorphism $\tau^*: U_i \rightarrow U_{n-i}$ allows us to identify m_{n-i} with $-m_i$, as it follows from the commutativity of the following diagram and the fact that $m_i \circ (\tau^* \otimes \text{Id}_{U_i}) = -m_i$

$$(17) \quad \begin{array}{ccc} \mathcal{L} \otimes U_i & \xrightarrow{m_i \circ (\tau^* \otimes \text{Id}_{U_i})} & U_i \\ \text{Id}_{\mathcal{L}} \otimes \tau^* \downarrow & & \downarrow \tau^* \\ \mathcal{L} \otimes U_{n-i} & \xrightarrow{m_{n-i}} & U_{n-i} \end{array}$$

3. For any $i, j = 1, \dots, n-1$, the product (7) induces, by restriction, a morphism

$$m_{ij}: U_i \otimes U_j \rightarrow U_{\overline{i+j}},$$

where $\overline{i+j} \in \{0, \dots, n-1\}$, $\overline{i+j} = i+j \pmod{n}$, and by definition $U_0 = q_*\mathcal{O}_Z$. In particular, for any $i = 1, \dots, n-1$, there is a morphism

$$m_{i,n-i}: U_i \otimes U_{n-i} \rightarrow \mathcal{O}_Y \oplus \mathcal{L}.$$

In the following, we will denote again with $m_{i,n-i}$ the previous morphism under the identification $\tau^*: U_{n-i} \rightarrow U_i$, hence

$$m_{i,n-i}: U_i \otimes U_i \rightarrow \mathcal{O}_Y \oplus \mathcal{L}.$$

Let $m_{i,n-i}^+: U_i \otimes U_i \rightarrow \mathcal{O}_Y$ and $m_{i,n-i}^-: U_i \otimes U_i \rightarrow \mathcal{L}$ be the compositions of $m_{i,n-i}$ with the projections onto \mathcal{O}_Y and \mathcal{L} respectively. Notice that, with respect to the involution on $U_i \otimes U_i$ that exchanges the factors, $m_{i,n-i}^+$ is symmetric, while $m_{i,n-i}^-$ is antisymmetric, hence they can be seen as morphisms

$$(18) \quad m_{i,n-i}^+: \text{Sym}^2(U_i) \rightarrow \mathcal{O}_Y, \quad m_{i,n-i}^-: \wedge^2(U_i) \rightarrow \mathcal{L},$$

or equivalently as sections

$$(19) \quad m_{i,n-i}^+ \in H^0(Y, \text{Sym}^2(U_i^\vee)), \quad m_{i,n-i}^- \in H^0(Y, \wedge^2(U_i^\vee) \otimes \mathcal{L}).$$

Proposition 5.5. *For any $i = 1, \dots, n-1$ the following statements hold true.*

- (i) $m_{i,n-i}^+$ is determined by m_i and $m_{i,n-i}^-$.
- (ii) The divisor of zeros of $m_{i,n-i}^-$ coincides with the divisor Δ_p defined in (16). In particular $m_{i,n-i}^-$ yields an isomorphism between $\wedge^2 U_i$ and $\mathcal{L} \otimes \mathcal{O}_Y(-\Delta_p)$.

Proof. (i) Let $y \in Y$, let $(U_i)_y$ be the stalk of U_i over y and let $s_1, s_2 \in (U_i)_y$. Then

$$m_{i,n-i}(s_1 \otimes s_2) = s_1 \tau^*(s_2),$$

hence

$$\begin{aligned} m_{i,n-i}^+(s_1 \otimes s_2) &= \frac{1}{2} (s_1 \tau^*(s_2) + \tau^*(s_1) s_2), \\ m_{i,n-i}^-(s_1 \otimes s_2) &= \frac{1}{2} (s_1 \tau^*(s_2) - \tau^*(s_1) s_2); \end{aligned}$$

where the product $s_1 \tau^*(s_2)$ (respectively $\tau^*(s_1) s_2$) is the usual one between stalks of regular functions defined in some neighborhood of $\pi^{-1}(y)$. For any $r \in (\mathcal{L})_y$, the associativity of the multiplication implies that

$$m_{i,n-i}(m_i(r \otimes s_1) \otimes s_2) = m(r \otimes m_{i,n-i}(s_1 \otimes s_2))$$

and hence

$$m_{i,n-i}^\pm(m_i(r \otimes s_1) \otimes s_2) = m(r \otimes m_{i,n-i}^\mp(s_1 \otimes s_2)).$$

This implies that, under the natural identification $\mathcal{O}_{Y,y} \cong \text{End}((\mathcal{L})_y)$, $m_{i,n-i}^+(s_1 \otimes s_2) = m_{i,n-i}^-(m_i((-) \otimes s_1) \otimes s_2)$, and hence the claim follows.

(ii) Without loss of generality we assume that Z is smooth. Indeed, since Z is normal, its singular locus $\text{Sing}(Z)$ has codimension ≥ 2 , so, for $Z^0 := Z \setminus \text{Sing}(Z)$ and $Y^0 := q(Z^0)$, the divisor of zeros of $m_{i,n-i}^-$ and Δ_p are determined by their restrictions to Y^0 .

For any $y \in Y$, let us consider the stalk $(m_{i,n-i}^-)_y$ of $m_{i,n-i}^-$ at y . Let u_i, v_i be a basis of $(U_i)_y$ as $\mathcal{O}_{Y,y}$ -module, then $u_i \otimes v_i - v_i \otimes u_i$ is a basis of $(\wedge^2 U_i)_y$, viewed as the submodule of $U_i \otimes U_i$. We have:

$$\begin{aligned} (20) \quad m_{i,n-i}^-(u_i \otimes v_i - v_i \otimes u_i) &= u_i \tau^*(v_i) - v_i \tau^*(u_i) \\ &= u_i \tau^*(v_i) - \tau^*(u_i \tau^*(v_i)), \end{aligned}$$

where we consider u_i and v_i as stalks of regular functions defined on some open neighbourhood of $\pi^{-1}(y)$ in X , such that $\sigma^*(u_i) = \exp(\frac{2\pi\sqrt{-1}}{n}i)u_i$ and $\sigma^*(v_i) = \exp(\frac{2\pi\sqrt{-1}}{n}i)v_i$.

Let us choose local analytic coordinates (y_1, \dots, y_d) for Y at y , and w on Z , such that Z is given locally by the equation $w^2 = y_1$. Furthermore, for any $i = 1, \dots, n-1$, let e_i be a basis of \mathcal{F}_i as $\left(\frac{\mathbb{C}[y_1, \dots, y_d, w]}{(w^2 - y_1)}\right)$ -module. Then let

$$(21) \quad u_i := 1 \cdot e_i, \quad v_i := w \cdot e_i.$$

Notice that we can choose the e_i 's in such a way that $\tau^*(e_i) = e_{n-i}$, for any $i = 1, \dots, n-1$, since τ^* identifies \mathcal{F}_i with \mathcal{F}_{n-i} . So, substituting (21) in (20) and using the equations $\tau^*(w) = -w$ and $\tau^*(e_i) = e_{n-i}$,

we obtain:

$$\begin{aligned}
m_{i,n-i}^-(u_i \otimes v_i - v_i \otimes u_i) &= u_i \tau^*(v_i) - \tau^*(u_i \tau^*(v_i)) \\
&= e_i \tau^*(we_i) - \tau^*(e_i \tau^*(we_i)) \\
&= e_i(-we_{n-i}) - \tau^*(e_i(-we_{n-i})) \\
&= -we_i e_{n-i} - we_i e_{n-i} \\
(22) \qquad \qquad \qquad &= -2we_i e_{n-i} = -2wb_p,
\end{aligned}$$

where we have used the fact that $e_i e_{n-i} = b_p$, b_p is a local equation for \mathcal{B}_p (this follows from (11)). Finally, let δ_p be a local equation for Δ_p such that $q^*(\delta_p) = b_p$, then the previous computation shows that

$$m_{i,n-i}^-(u_i \otimes v_i - v_i \otimes u_i) = -2wq^*(\delta_p),$$

from which the claim follows. \square

The following result is a converse to Proposition 5.3.

Theorem 5.6 (Structure of D_n -covers). *Let Y be a smooth variety and n be a positive integer. Then, to the following data a), b) and c), we can associate a D_n -cover $\pi: X \rightarrow Y$ in a natural way.*

- a) A line bundle \mathcal{L} and an effective reduced divisor \mathcal{B}_q on Y , such that $\mathcal{L}^{\otimes 2} \cong \mathcal{O}_Y(-\mathcal{B}_q)$.
- b) Reduced effective Weil divisors $D_1, \dots, D_{\lfloor \frac{n}{2} \rfloor}$ on $Z := \text{Spec}(\mathcal{O}_Y \oplus \mathcal{L})$ without common components, such that $\bar{\tau}(D_1 \cup \dots \cup D_{\lfloor \frac{n-1}{2} \rfloor})$ doesn't have common components with $D_1 \cup \dots \cup D_{\lfloor \frac{n-1}{2} \rfloor}$, and, in the case where n is even, $\bar{\tau}(D_{\frac{n}{2}}) = D_{\frac{n}{2}}$; where $\lfloor a \rfloor$ denotes the integer part of a number $a \in \mathbb{Q}$, and $\bar{\tau}$ is the involution of the double cover $q: Z \rightarrow Y$.
- c) Divisorial sheaves $\mathcal{F}_1, \dots, \mathcal{F}_{\lfloor \frac{n}{2} \rfloor}$ on Z flat over \mathcal{O}_Y , such that, for any $i, j = 1, \dots, n-1$, (13) holds, and if n is even $\mathcal{F}_{\frac{n}{2}} = \bar{\tau}^*(\mathcal{F}_{\frac{n}{2}})$; where for $\lfloor \frac{n}{2} \rfloor < \ell, k \leq n-1$, $\mathcal{F}_\ell := \bar{\tau}^*(\mathcal{F}_{n-\ell})$ and $D_k := \bar{\tau}(D_{n-k})$, the coefficients ε_{ij}^k are defined in (12).

The variety X so constructed is normal if and only if, setting $\varkappa := \gcd\{k = 1, \dots, n-1 \mid D_k \neq 0\}$, then either $\varkappa = 1$, or $\frac{n}{\varkappa} \mathcal{F}_1 - \sum_{k=1}^{n-1} \frac{k}{\varkappa} D_k$ has order precisely \varkappa in the group of divisorial sheaves of Z . In this case $\pi_* \mathcal{O}_X = \mathcal{O}_Y \oplus \mathcal{L} \oplus_{i=1}^{\lfloor \frac{n}{2} \rfloor} q_* \mathcal{F}_i$, in particular π is flat.

Before giving the proof of the previous theorem, two remarks are in order.

Remark 5.7. 1. Using the results of Section 3, the divisorial sheaves in c) correspond to pairs $(U_1, m_1), \dots, (U_{\lfloor \frac{n}{2} \rfloor}, m_{\lfloor \frac{n}{2} \rfloor})$ consisting of rank 2 vector bundles on Y , $U_1, \dots, U_{\lfloor \frac{n}{2} \rfloor}$, and morphisms $m_i: U_i \otimes \mathcal{L} \rightarrow U_i$ such that $m_i^2 = \delta_q \text{Id}_{U_i}$, where $\mathcal{B}_q = \{\delta_q = 0\}$. With this notation, $\bar{\tau}^* \mathcal{F}_i$ corresponds to $(U_i, -m_i)$, for any $i = 1, \dots, n-1$. Furthermore, using Proposition 3.3, the relation (13) can be written in terms of the (U_i, m_i) 's and the D_i 's.

2. A similar criterion for the existence of D_n -covers $\pi: X \rightarrow Y$ has been given in [Tok94], where X is constructed as the normalization of Y in a certain dihedral field extension of $\mathbb{C}(Y)$. Here, using the structure theorem for cyclic covers, we construct $\pi: X \rightarrow Y$ explicitly, we hope that this procedure could be useful for further investigations of Galois covers.

Proof. The data a) determines a flat double cover $q: Z \rightarrow Y$ in the usual way, $Z := \text{Spec}(\mathcal{O}_Y \oplus \mathcal{L})$ and q is given by the inclusion $\mathcal{O}_Y \hookrightarrow \mathcal{O}_Y \oplus \mathcal{L}$ as the first summand. Notice that Z is normal since \mathcal{B}_q is reduced.

We define $X := \text{Spec}(\mathcal{O}_Z \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_{n-1})$, where $\mathcal{O}_Z \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_{n-1}$ is the sheaf of \mathcal{O}_Z -algebras with algebra structure given by the isomorphisms (13) in the following way. Since $\mathcal{F}_{\overline{i+j}}$ is a divisorial sheaf, for $i, j = 1, \dots, n-1$, a morphism $\mathcal{F}_i \otimes \mathcal{F}_j \rightarrow \mathcal{F}_{\overline{i+j}}$ is uniquely determined by its restriction on the smooth locus Z^0 of Z . Let us fix firstly sections $\delta_k \in H^0(Z, \mathcal{O}_Z(D_k))$, such that $D_k = \{\delta_k = 0\}$ on Z^0 , for any $k = 1, \dots, n-1$. By Lemma 3.1, on Z^0 the sheaves $\mathcal{F}_1, \dots, \mathcal{F}_{n-1}$ are locally free, so, locally where they are trivial, we choose generators e_1, \dots, e_{n-1} such that $e_{n-i} = \overline{\tau}^*(e_i)$, for any i . The algebra structure on the restriction of $\mathcal{O}_Z \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_{n-1}$ on such open subsets is defined as usual by the equations

$$e_i e_j = e_{\overline{i+j}} \prod_{k=1}^{n-1} \delta_k^{\varepsilon_{ij}^k}, \quad \text{for any } i, j = 1, \dots, n-1.$$

Notice that, if we choose different generators, $\tilde{e}_1, \dots, \tilde{e}_{n-1}$ satisfying the same conditions $\tilde{e}_{n-i} = \overline{\tau}^*(\tilde{e}_i)$, then we obtain an algebra canonically isomorphic to the previous one. Hence the construction globalizes and $\mathcal{O}_Z \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_{n-1}$ becomes a sheaf of \mathcal{O}_Z -algebras. The morphism $\pi: X \rightarrow Y$ is defined as usual.

From the previous construction and from the fact that $D_{n-k} = \overline{\tau}(D_k)$, it follows that $\overline{\tau}^*: \mathcal{O}_Z \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_{n-1} \rightarrow \mathcal{O}_Z \oplus \mathcal{F}_1 \oplus \dots \oplus \mathcal{F}_{n-1}$ is a morphism of \mathcal{O}_Z -algebras. This defines an involution $\tau: X \rightarrow X$. By construction we have that $\sigma^* \circ \tau^* = \tau^* \circ (\sigma^{-1})^*$, hence X carries an action of D_n such that $\pi: X \rightarrow Y$ is a D_n -cover.

The condition for X to be normal follows from Theorem 1.1 in [Cat10]. \square

Remark 5.8. Similarly as for cyclic covers, X in Theorem 5.6 is determined by the data a), b) and \mathcal{F}_1 (or (U_1, m_1)), such that

$$(\mathcal{F}_1^{\otimes n})^{**} \cong \mathcal{O}_Z \left(- \sum_{k=1}^{n-1} k D_k \right).$$

Furthermore, by Proposition 4.4, $(\mathcal{F}_1^{\otimes n})^{**} = (\text{Sym}^n(U_1)) / \mathcal{K}_n$.

5.3. D_3 -covers and triple covers. In this section we restrict ourselves to D_3 -covers and relate the results that we obtained so far with the structure theorem for flat finite morphisms of degree 3 of algebraic varieties ([Mir85]), also called *triple covers*. This relation originates from the following fact: for every D_3 -cover $\pi: X \rightarrow Y$, any quotient $W := X/\langle\sigma^i\tau\rangle$ has a natural structure of triple cover of Y induced by π ; conversely, if $W \rightarrow Y$ is a triple cover which is not Galois, then its Galois closure is a D_3 -cover. Since the elements $\sigma^i\tau$ are pairwise conjugate, the corresponding triple covers are isomorphic, hence in the following we consider only $X/\langle\tau\rangle$. Notice that the structure of D_3 -covers has been studied also in [East11].

Let $\pi: X \rightarrow Y$ be a D_3 -cover. We have seen in the previous section that π is determined by the locally free sheaves \mathcal{L}, U_1, U_2 and a morphism of \mathcal{O}_Y -modules

$$m: (\mathcal{O}_Y \oplus \mathcal{L} \oplus U_1 \oplus U_2)^{\otimes 2} \rightarrow \mathcal{O}_Y \oplus \mathcal{L} \oplus U_1 \oplus U_2$$

which gives an associative commutative product. The involution τ yields an isomorphism $\tau^*: U_1 \rightarrow U_2$ as explained in the previous section. Under this identification one easily see that $\pi: X \rightarrow Y$ is determined by $\mathcal{O}_Y, \mathcal{L}, U_1, m_{12}, m_1$ and m_{11} (see the previous section for their definitions). In particular,

$$\mathcal{O}_X = \mathcal{O}_Y \oplus \mathcal{L} \oplus U_1 \oplus U_1$$

and the involution τ^* acts on the element $(a, b, c, d) \in \mathcal{O}_Y \oplus \mathcal{L} \oplus U_1 \oplus U_1$ as follows:

$$\tau^*(a, b, c, d) = (a, -b, d, c).$$

The subsheaf of τ^* -invariants, $(\mathcal{O}_X)^{\tau^*} \subset \mathcal{O}_X$, consists of the elements of the form $(a, 0, c, c) \in \mathcal{O}_Y \oplus \mathcal{L} \oplus U_1 \oplus U_1$, so the sheaf of regular functions on $W := X/\langle\tau\rangle$ is

$$(23) \quad \mathcal{O}_W \cong \mathcal{O}_Y \oplus U_1$$

with inclusion in $\mathcal{O}_Y \oplus \mathcal{L} \oplus U_1 \oplus U_1$ given by $(a, c) \mapsto (a, 0, c, c)$. The morphism $\pi: X \rightarrow Y$ descends to a morphism $f: W \rightarrow Y$, which is a triple cover. We refer to [Mir85] for definitions, notations and results concerning triple covers.

Proposition 5.9. *Under the identification (23), U_1 coincides with the Tschirnhausen module of \mathcal{O}_W over \mathcal{O}_Y , that is U_1 consists of the elements of \mathcal{O}_W whose minimal cubic polynomial has no square term ($U_1 = E$ in the notation of [Mir85]). Furthermore, the tensor ϕ_2 in [Mir85] coincides with $m_{11}: U_1 \otimes U_1 \rightarrow U_1$, which has been defined in the previous section.*

Proof. For $(0, c) \in \mathcal{O}_Y \oplus U_1$, we calculate its cubic power $(0, c)^3$. To this aim, we consider its image $(0, 0, c, c) \in \mathcal{O}_Y \oplus \mathcal{L} \oplus U_1 \oplus U_1$ and use the product m of \mathcal{O}_X . In order to simplify the notation, we write

$(0, 0, c, c) = c + \tau^*(c) \in \mathcal{O}_X$ and denote the product m simply by \cdot . Then we have the following expression:

$$(c + \tau^*(c))^3 = c^3 + \tau^*(c)^3 + 2c\tau^*(c)(c + \tau^*(c)).$$

Since $c^3 + \tau^*(c)^3, 2c\tau^*(c) \in \mathcal{O}_Y$, the minimal cubic polynomial of $c + \tau^*(c)$ has no square term and so it belongs to the Tschirnhausen module of \mathcal{O}_W over \mathcal{O}_Y . This proves the first claim.

For the second claim, recall that by definition ϕ_2 is the composition of the product in \mathcal{O}_W followed by the projection onto U_1 . For any $c, c' \in U_1$, the product between $c + \tau^*(c), c' + \tau^*(c') \in \mathcal{O}_W$ has the following expression:

$$(c + \tau^*(c))(c' + \tau^*(c')) = cc' + \tau^*(cc') + c\tau^*(c') + \tau^*(c\tau^*(c')).$$

Notice that $c\tau^*(c') + \tau^*(c\tau^*(c')) \in \mathcal{O}_Y$ and that $cc' + \tau^*(cc')$ corresponds to $(0, \tau^*(cc')) \in U_1$ under the identification (23). So ϕ_2 coincides with m_{11} under the identification $\tau^*: U_1 \rightarrow U_2$. \square

Let $f: W \rightarrow Y$ be a flat finite map of degree 3, let E be the Tschirnhausen module of \mathcal{O}_W over \mathcal{O}_Y and let $\phi_2: \text{Sym}^2(E) \rightarrow E$ be the associated triple cover homomorphism, as defined in [Mir85]. Under the standard identification $E \cong E^\vee \otimes \wedge^2(E) = \text{Hom}(E, \wedge^2(E))$, $u \mapsto (v \mapsto u \wedge v)$, we can view ϕ_2 as a morphism $\phi_2: \text{Sym}^2(E) \otimes E \rightarrow \wedge^2(E)$. By [Mir85, Prop. 3.5], ϕ_2 is symmetric, hence it induces a morphism $\Phi: \text{Sym}^3(E) \rightarrow \wedge^2(E)$. The structure theorem for triple covers states that conversely $f: W \rightarrow Y$ is determined by a rank 2 locally free sheaf E of \mathcal{O}_Y -modules and an \mathcal{O}_Y -morphism $\Phi: \text{Sym}^3(E) \rightarrow \wedge^2(E)$ ([Mir85, Thm. 3.6]). The proof of the fact that ϕ_2 is symmetric given in [Mir85] is done by a direct calculation using local coordinates and is a consequence of the fact that E is the Tschirnhausen module of \mathcal{O}_W over \mathcal{O}_Y .

We extend part of these results to D_n -covers, $n \geq 2$, and give a different proof of [Mir85, Prop. 3.5] quoted above. For any D_n -cover $\pi: X \rightarrow Y$, there is a \mathcal{O}_Y -morphism

$$\phi_{n-1}: \text{Sym}^{n-1}(U_1) \rightarrow U_1$$

defined as follows: for any $u_1 \otimes \dots \otimes u_{n-1} \in \text{Sym}^{n-1}(U_1)$, their product $m(u_1 \otimes \dots \otimes u_{n-1})$ is in U_{n-1} , so applying $\tau^*: U_{n-1} \rightarrow U_1$ we obtain an element in U_1 , this is by definition $\phi_{n-1}(u_1 \otimes \dots \otimes u_{n-1})$. Under the identification $U_1 \cong U_1^\vee \otimes \wedge^2(U_1)$ as before, we can see ϕ_{n-1} as a morphism

$$\phi_{n-1}: \text{Sym}^{n-1}(U_1) \otimes U_1 \rightarrow \wedge^2(U_1).$$

Proposition 5.10. *The morphism ϕ_{n-1} is symmetric, hence induces a morphism of \mathcal{O}_Y -modules $\Phi_{n-1}: \text{Sym}^n(U_1) \rightarrow \wedge^2(U_1)$.*

Proof. By the commutativity of m , Φ_{n-1} is symmetric in the first $n-1$ entries. So, it is enough to prove that

$$(24) \quad \Phi_{n-1}(u_1 \otimes \dots \otimes u_{n-1} \otimes v) = \Phi_{n-1}(u_1 \otimes \dots \otimes u_{n-2} \otimes v \otimes u_{n-1}),$$

for any $u_1, \dots, u_{n-1}, v \in U_1$. To this aim, let us consider the morphism

$$m_{1,n-1}^-: \Lambda^2(U_1) \rightarrow \mathcal{L}$$

defined in (18). By Proposition 5.5, $m_{1,n-1}^-$ identifies $\Lambda^2(U_1)$ with $\mathcal{L}(-\Delta_p)$, hence (24) is equivalent to

$$(25) \quad \begin{aligned} m_{1,n-1}^- (\Phi_{n-1}(u_1 \otimes \dots \otimes u_{n-1} \otimes v)) &= \\ &= m_{1,n-1}^- (\Phi_{n-1}(u_1 \otimes \dots \otimes u_{n-2} \otimes v \otimes u_{n-1})) . \end{aligned}$$

From the explicit form of the isomorphism $U_1 \cong U_1^\vee \otimes \Lambda^2(U_1)$ recalled before, we have that

$$\Phi_{n-1}(u_1 \otimes \dots \otimes u_{n-1} \otimes v) = \tau^*(m(u_1 \otimes \dots \otimes u_{n-1})) \wedge v .$$

Furthermore, by definition, $m_{1,n-1}^- (\tau^*(m(u_1 \otimes \dots \otimes u_{n-1})) \wedge v)$ is the projection onto \mathcal{L} of $m(\tau^*(m(u_1 \otimes \dots \otimes u_{n-1})) \otimes \tau^*(v)) \in \mathcal{O}_Y \oplus \mathcal{L}$. Since τ^* commutes with m ,

$$m(\tau^*(m(u_1 \otimes \dots \otimes u_{n-1})) \otimes \tau^*(v)) = \tau^*m(m(u_1 \otimes \dots \otimes u_{n-1})) \otimes v .$$

Using the associativity and the commutativity of m , we deduce that

$$\tau^*m(m(u_1 \otimes \dots \otimes u_{n-1})) \otimes v = \tau^*m(m(u_1 \otimes \dots \otimes u_{n-2} \otimes v)) \otimes u_{n-1} .$$

Projecting both sides of this equation onto \mathcal{L} we obtain (25), from which the statement follows. \square

Remark 5.11. From Theorem 5.6 it follows that, for a D_n -cover $\pi: X \rightarrow Y$, the intermediate degree n cover $f: W := X/\langle \tau \rangle \rightarrow Y$ embeds naturally in $\mathcal{O}_Y \oplus U_1 \oplus \dots \oplus U_{\frac{n-1}{2}}$ if n is odd, respectively in $\mathcal{O}_Y \oplus U_1 \oplus \dots \oplus U_{\frac{n}{2}-1} \oplus \mathcal{M}$ when n is even. Then the symmetric morphism $\Phi_{n-1}: \text{Sym}^n(U_1) \rightarrow \Lambda^2(U_1)$ determines $f: W \rightarrow Y$ birationally.

6. SIMPLE AND ALMOST SIMPLE DIHEDRAL COVERS AND THEIR INVARIANTS

In this section we define the *simple dihedral covers* and the *almost simple dihedral covers*, and we investigate some of their properties. Throughout the section we assume that Y is a smooth variety.

Recall that, from Proposition 5.2, a D_n -field extension $\mathbb{C}(Y) \subset E$ is given by $E = \mathbb{C}(Y)(x)$, where x satisfies an irreducible equation of the following form:

$$x^{2n} - 2ax^n + F^n = 0 ,$$

with $a, F \in \mathbb{C}(Y)$. In order to construct varieties having E as field of rational functions, one could proceed in the following way. Consider a geometric line bundle $\mathbb{L} \rightarrow Y$, sections $a \in H^0(Y, \mathbb{L}^{\otimes n})$, $F \in$

$H^0(Y, \mathbb{L}^{\otimes 2})$, and define

$$X' := \{v \in \mathbb{L} \mid v^{2n} - 2av^n + F^n = 0\} \subset \mathbb{L},$$

where v is a fibre coordinate of \mathbb{L} . The singular locus of X' contains the locus $v = F = 0$, so it has codimension 1 and X' is not normal.

We slightly modify this construction in the following way. Let $\mathbb{L} \rightarrow Y$ be a geometric line bundle, $a \in H^0(Y, \mathbb{L}^{\otimes n})$ and $F \in H^0(Y, \mathbb{L}^{\otimes 2})$. Define $X \subset \mathbb{L} \oplus \mathbb{L}$ to be the set of $(u, v) \in \mathbb{L} \oplus \mathbb{L}$ such that the following equations are satisfied:

$$(26) \quad \begin{cases} uv & = F \\ u^n - 2a + v^n & = 0, \end{cases}$$

where u (resp. v) is a fibre coordinate of the first (resp. second) copy of \mathbb{L} in $\mathbb{L} \oplus \mathbb{L}$. The dihedral group D_n acts on X via $\sigma(u, v) = (\zeta u, \zeta^{-1}v)$, $\tau(u, v) = (v, u)$, where $\zeta \in \mathbb{C}^*$ is a primitive root of 1.

Theorem 6.1. *Let Y be a smooth variety. Let $\mathbb{L} \rightarrow Y$ be a geometric line bundle, $a \in H^0(Y, \mathbb{L}^{\otimes n})$ and $F \in H^0(Y, \mathbb{L}^{\otimes 2})$, such that:*

- (i) *the zero locus of $a^2 - F^n \in H^0(Y, \mathbb{L}^{\otimes 2n})$ is smooth in the open set $F \neq 0$;*
- (ii) *the divisors $\{a = 0\}$ and $\{F = 0\}$ intersect each other transversely.*

Then the variety X defined by the equations (26) is smooth and the restriction to X of the fibre bundle projection $\mathbb{L} \oplus \mathbb{L} \rightarrow Y$ is a D_n -cover, $\pi: X \rightarrow Y$, with branch divisor $\mathcal{B}_\pi = \{F^n - a^2 = 0\}$. Furthermore, if $\{a = 0\} \cap \{F = 0\} \neq \emptyset$, then X is irreducible.

Definition 6.2. *A simple D_n -cover of Y is the D_n -cover $\pi: X \rightarrow Y$ given as in Theorem 6.1 by the restriction to X of the fibre bundle projection $\mathbb{L} \oplus \mathbb{L} \rightarrow Y$.*

Proof. (Of Theorem 6.1.) Let us define $\Phi_1 := uv - F$ and $\Phi_2 := u^n - 2a + v^n$, so that the equations (26) become

$$\begin{cases} \Phi_1 & = 0 \\ \Phi_2 & = 0. \end{cases}$$

Taking partial derivatives along the fibre coordinates we have:

$$\begin{pmatrix} \frac{\partial \Phi_1}{\partial u} & \frac{\partial \Phi_1}{\partial v} \\ \frac{\partial \Phi_2}{\partial u} & \frac{\partial \Phi_2}{\partial v} \end{pmatrix} = \begin{pmatrix} v & u \\ nu^{n-1} & nv^{n-1} \end{pmatrix},$$

hence the ramification divisor of π is $R = \{v^n - u^n = 0\}$. Notice that $v^n - u^n$ is $\langle \sigma^* \rangle$ -invariant, while $\tau^*(v^n - u^n) = -(v^n - u^n)$. Therefore

the branch locus \mathcal{B}_π is defined by the following equation:

$$\begin{aligned} (v^n - u^n)(u^n - v^n) &= -(v^n - u^n)^2 \\ &= -(v^n + u^n)^2 + 4(uv)^n \\ &= -4a^2 + 4F^n. \end{aligned}$$

Let us consider first the restriction of π over the points where $F \neq 0$. In this locus, we have $v = F/u$ and so the equation $\Phi_2 = 0$ is equivalent to

$$\Phi = 0, \quad \text{where} \quad \Phi := u^{2n} - 2au^n + F^n.$$

It follows that π is finite over the open set $F \neq 0$. To show the flatness of π over $F \neq 0$, by Prop. 2.2 it suffices to prove that X is smooth there. To this aim consider

$$\frac{\partial \Phi}{\partial u} = 2nu^{n-1}(u^n - a),$$

which can vanish only if $u^n - a = 0$, since $F \neq 0 \Rightarrow u \neq 0$. On the other hand, over the locus $u^n - a = 0$,

$$\begin{aligned} \frac{\partial \Phi}{\partial y} &= -2u^n \frac{\partial a}{\partial y} + \frac{\partial F^n}{\partial y} \\ &= \frac{\partial (F^n - a^2)}{\partial y}, \end{aligned}$$

where y is any coordinate function on Y . Since $F^n - a^2 = 0$ is the branch divisor and, by hypothesis (i), it is smooth in $F \neq 0$, it follows that X is smooth there.

We consider now the restriction of π over the locus $F = 0$ and notice that here $uv = 0$. If $u \neq 0$, then as before we see that π is finite. The smoothness of X at $(u \neq 0, v = 0)$ follows from the fact that $(u \neq 0, v = 0) \notin R$. The same argument applies if $v \neq 0$. It remains the case where $u = v = 0$, which implies that $a = F = 0$. By hypothesis (ii), a and F are part of a local coordinate system near the points where $a = F = 0$, in the analytic topology. Writing the equations for X in these coordinates, we see that X is smooth at these points. There remains to show that π is finite also over the points $a = F = 0$. Here uv and $u^n + v^n$ are precisely the D_n -invariants, and \mathcal{O}_X is a free \mathcal{O}_Y -module generated by $1, u^n - v^n, u, u^2, \dots, u^{n-1}, v, v^2, \dots, v^{n-1}$.

Let us now assume that $\{a = 0\} \cap \{F = 0\} \neq \emptyset$. The irreducibility of X is equivalent to the surjectivity of the monodromy of the cover, $\mu: \pi_1(Y \setminus \mathcal{B}_\pi) \rightarrow D_n$. Since the branch divisor of the intermediate cover $q: Z \rightarrow Y$ coincides with \mathcal{B}_π , it is reduced, so Z is irreducible and hence $\text{Im}(\mu)$ contains a reflection $\sigma^i \tau$. To see that also $\sigma \in \text{Im}(\mu)$, let us choose local coordinates $(a, F, y_3, \dots, y_{\dim(Y)})$ for Y at a point in $\{a = 0\} \cap \{F = 0\}$ and consider the path in X

$$\gamma(t) = \left(\exp\left(\frac{2\pi\sqrt{-1}}{n}t\right) u_0, \exp\left(-\frac{2\pi\sqrt{-1}}{n}t\right) v_0, F, a, y_3, \dots, y_{\dim(Y)} \right),$$

where, $(u_0, v_0, F, a, y_3, \dots, y_{\dim(Y)}) \in X$. Then, $\gamma(1) = \sigma\gamma(0)$, and $\pi \circ \gamma$ is a loop contained in $Y \setminus \mathcal{B}_\pi$, if and only if

$$\exp(2\pi\sqrt{-1}t)u_0^n - \exp(-2\pi\sqrt{-1}t)v_0^n \neq 0, \quad \forall t.$$

Since this condition can be easily achieved, e.g. by choosing u_0, v_0 with $|u_0| \neq |v_0|$, the claim follows. \square

Notice that the hypothesis (i) in the previous theorem is general, as it follows from the next proposition.

Proposition 6.3. *Under the same notation as before, for a general choice of $a \in H^0(Y, \mathbb{L}^{\otimes n})$ and $F \in H^0(Y, \mathbb{L}^{\otimes 2})$, $F^n - a^2 = 0$ is smooth over $F \neq 0$.*

Proof. This follows directly from Bertini's theorem (see e.g. [GH]). \square

6.1. Invariants of simple dihedral covers. We first determine the eigensheaves decomposition of $\pi_*\mathcal{O}_X$, where $\pi: X \rightarrow Y$ is a simple D_n -cover. Let \mathbb{L}, F and a be as in the statement of Thm. 6.1, and let us denote with $\mathcal{O}_Y(L)$ the sheaf of sections of \mathbb{L} . Then, over an open subset $S \subset Y$ where \mathbb{L} is trivial, from (26) we deduce that

$$\begin{aligned} \pi_*\mathcal{O}_{X|S} &\cong \frac{\mathcal{O}_S[\lambda, \mu]}{(\lambda\mu - F, \lambda^n - 2a + \mu^n)} \\ &\cong \mathcal{O}_S \oplus \mathcal{O}_S(\lambda^n - \mu^n) \oplus_{i=1}^{n-1} (\mathcal{O}_S\lambda^i \oplus \mathcal{O}_S\mu^{n-i}), \end{aligned}$$

where λ and μ are fibre coordinates on the dual \mathbb{L}^\vee of \mathbb{L} . This local description globalizes and we have:

$$(27) \quad \pi_*\mathcal{O}_X \cong \bigoplus_{i=0}^{n-1} [\mathcal{O}_Y(-iL) \oplus \mathcal{O}_Y(-(n-i)L)].$$

Notice that, for the sheaves \mathcal{L} and U_i introduced in Section 5.2, we have:

$$\mathcal{L} = \mathcal{O}_Y(-nL), \quad U_i = \mathcal{O}_Y(-iL) \oplus \mathcal{O}_Y(-(n-i)L).$$

To determine the canonical bundle of X , we use the canonical bundle formula for branched covers:

$$\omega_X = \pi^*\omega_Y \otimes \mathcal{O}_X(R),$$

where R is the ramification divisor of π . If $\pi: X \rightarrow Y$ is a simple D_n -cover, then $R = \{u^n - v^n = 0\}$ (see the proof of Thm. 6.1). Since $u^n - v^n$ is a generator of the eigensheaf corresponding to the irreducible representation of D_n with character χ_2 , $\mathcal{O}_X(R) = \pi^*\mathcal{L}^\vee = \pi^*\mathcal{O}_Y(nL)$. Hence

$$(28) \quad \omega_X = \pi^*(\omega_Y(nL)).$$

In particular, if $\dim(X) = 2$, then the self-intersection of a canonical divisor of X is:

$$(29) \quad K_X^2 = 2n(K_Y + nL)^2.$$

To compute the Euler characteristic $\chi(\mathcal{O}_X)$, by the finiteness of π we have that $\chi(\mathcal{O}_X) = \chi(\pi_*\mathcal{O}_X)$. In particular, if $\dim(X) = 2$, the Riemann-Roch theorem for surfaces yields the following formula:

$$(30) \quad \chi(\mathcal{O}_X) = 2n\chi(\mathcal{O}_Y) + \frac{1}{6}n(2n^2 + 1)L \cdot L + \frac{1}{2}n^2L \cdot K_Y.$$

6.2. Almost simple dihedral covers. In this section we define the almost simple dihedral covers, which can be seen as projectivizations of the simple dihedral covers. The construction follows closely the one of almost simple cyclic covers, introduced and studied in [Cat89].

The covering space X of an almost simple dihedral cover $\pi: X \rightarrow Y$, over the smooth variety Y , is defined as a complete intersection in the fibre product $\mathbb{P}_1 \times_Y \mathbb{P}_2 \rightarrow Y$ of two \mathbb{P}^1 -bundles over Y in the following way. Let $\mathbb{L} \rightarrow Y$ be a geometric line bundle and $\underline{\mathbb{C}} = Y \times \mathbb{C} \rightarrow Y$ be the trivial geometric line bundle. For each $i = 1, 2$, let $\mathbb{P}_i := \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L})$ be the \mathbb{P}^1 -bundle associated to $\underline{\mathbb{C}} \oplus \mathbb{L} \rightarrow Y$, and let $p_i: \mathbb{P}_i \rightarrow Y$ be the corresponding projection. Here, each fibre of p_i is the projective space of 1-dimensional subspaces in the corresponding fibre of $\underline{\mathbb{C}} \oplus \mathbb{L}$, hence, using the notation in [Hart77], $\mathbb{P}_i = \text{Proj}(\mathcal{O}_Y \oplus \mathcal{O}_Y(-L))$ and in particular $(p_i)_*\mathcal{O}_{\mathbb{P}_i}(1) = \mathcal{O}_Y \oplus \mathcal{O}_Y(-L)$, where L is a divisor in Y such that $\mathcal{O}_Y(L)$ is the sheaf of sections of \mathbb{L} (hence $\mathbb{L} = \text{Spec}(\text{Sym}(\mathcal{O}_Y(-L)))$). On each \mathbb{P}_i there are projective fibre coordinates: $[u_0 : u_1]$ for \mathbb{P}_1 , $[v_0 : v_1]$ for \mathbb{P}_2 . u_0, u_1 are defined as follows (v_0, v_1 are defined in the same way): let $U \subset p_1^*(\underline{\mathbb{C}} \oplus \mathbb{L})$ be the universal sub-bundle, then $u_0: U \rightarrow p_1^*(\underline{\mathbb{C}})$ is the composition of the inclusion with the projection; similarly for $u_1: U \rightarrow p_1^*\mathbb{L}$. Notice that, since $U = \text{Spec}(\text{Sym } \mathcal{O}_{\mathbb{P}_1}(1))$, then $u_0 \in H^0(\mathbb{P}_1, \mathcal{O}_{\mathbb{P}_1}(1))$ and $u_1 \in H^0(\mathbb{P}_1, \mathcal{O}_{\mathbb{P}_1}(1) \otimes p_1^*\mathcal{O}_Y(L))$.

Let $F \in H^0(Y, \mathbb{L}^{\otimes 2})$ and let A_0, A_∞ be effective divisors in Y such that

$$A_0 \equiv nL + A_\infty.$$

Let $a_\infty \in H^0(Y, \mathcal{O}_Y(A_\infty))$ and $a_0 \in H^0(Y, \mathcal{O}_Y(A_0))$ be such that $A_\infty = \{a_\infty = 0\}$ and $A_0 = \{a_0 = 0\}$.

Consider the subvariety $X \subset \mathbb{P}_1 \times_Y \mathbb{P}_2 = \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \times_Y \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L})$ defined by the following equations:

$$(31) \quad \begin{cases} \Phi_1 &= 0 \\ \Phi_2 &= 0, \end{cases}$$

where

$$\Phi_1 := u_1v_1 - u_0v_0F, \quad \Phi_2 := a_\infty v_1^n u_0^n - 2a_0 v_0^n u_0^n + a_\infty v_0^n u_1^n,$$

u_0, u_1, v_0, v_1 are defined above and $([u_0 : u_1], [v_0 : v_1])$ are projective fibre coordinates on $\mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \times_Y \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L})$.

Notice that the dihedral group D_n acts on X in the following way:

$$\begin{aligned} \sigma([u_0 : u_1], [v_0 : v_1]) &= ([u_0 : \zeta u_1], [v_0 : \zeta^{-1}v_1]), \\ \tau([u_0 : u_1], [v_0 : v_1]) &= ([v_0 : v_1], [u_0 : u_1]), \end{aligned}$$

where $\zeta \in \mathbb{C}^*$ is a primitive n -th root of 1. Then we have the following result.

Theorem 6.4. *Let Y be a smooth variety, $\mathbb{L} \rightarrow Y$ be a geometric line bundle, $\underline{\mathbb{C}} = Y \times \mathbb{C} \rightarrow Y$ be the trivial geometric line bundle, and $F \in H^0(Y, \mathbb{L}^{\otimes 2})$. Let A_0, A_∞ be effective divisors in Y such that*

$$A_0 \equiv nL + A_\infty,$$

where L is a divisor in Y with $\mathcal{O}_Y(L)$ being the sheaf of sections of \mathbb{L} . Let $a_\infty \in H^0(Y, \mathcal{O}_Y(A_\infty))$ and $a_0 \in H^0(Y, \mathcal{O}_Y(A_0))$ be such that $A_\infty = \{a_\infty = 0\}$ and $A_0 = \{a_0 = 0\}$.

Assume that the following conditions are satisfied:

- (i) A_0 intersects $\{F = 0\}$ transversely, $A_0 \cap A_\infty = \emptyset$ and A_∞ is smooth;
- (ii) the locus $\{a_0^2 - F^n a_\infty^2 = 0\}$ is smooth on the open set $F \neq 0$.

Then X , defined by (31), is smooth and the restriction of the projection $\mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \times_Y \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \rightarrow Y$ to X is a D_n -cover $\pi: X \rightarrow Y$ with branch divisor $\mathcal{B}_\pi = \{a_\infty(a_0^2 - a_\infty^2 F^n) = 0\}$. Furthermore, if $A_0 \cap \{F = 0\} \neq \emptyset$, then X is irreducible.

Definition 6.5. *An almost simple D_n -cover of Y is the D_n -cover $\pi: X \rightarrow Y$ given as in Theorem 6.4 by the restriction to X of the fibre bundle projection $\mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \times_Y \mathbb{P}(\underline{\mathbb{C}} \oplus \mathbb{L}) \rightarrow Y$.*

Proof. We first prove that the intersection of X with each one of the standard open subsets $v_0 u_0 \neq 0$, $v_1 u_0 \neq 0$ and $v_1 u_1 \neq 0$ is a smooth variety. This suffices because $\tau(\{v_0 u_1 \neq 0\}) = \{v_1 u_0 \neq 0\}$.

To this aim, observe that the restriction of π to the locus where $u_0 v_0 \neq 0$ is a simple D_n -cover. Indeed, setting $u = u_1/u_0$ and $v = v_1/v_0$, the equations (31) reduce to

$$\begin{cases} uv - F & = 0 \\ a_\infty v^n - 2a_0 + a_\infty u^n & = 0. \end{cases}$$

Since $A_\infty \cap A_0 = \emptyset$, on this locus a_∞ never vanishes, and setting $a = a_0/a_\infty$ we obtain the equations (26). Under our hypotheses Theorem 6.1 applies, hence X is smooth if $u_0 v_0 \neq 0$.

Consider now the locus $v_1 u_1 \neq 0$ and observe that there F never vanishes. Let us define $u = u_0/u_1$, $v = v_0/v_1$ and $g = 1/F$. Then the equations (31) become

$$\begin{cases} uv & = g \\ a_\infty u^n - 2a_0 g^n + a_\infty v^n & = 0. \end{cases}$$

Since $A_\infty \cap A_0 = \emptyset$, a_∞ never vanishes in this locus. So, defining $a = a_0/a_\infty$, we get the following equations for X :

$$u^{2n} - 2ag^n u^n + g^n = 0, \quad v = g/u.$$

Notice that this is the equation of a D_n -cover, with action $\sigma(u) = \zeta u$, $\tau(u) = g/u$. Since $g \neq 0$ everywhere, $\langle \sigma \rangle \cong \mathbb{Z}/n\mathbb{Z}$ acts freely and so X is smooth if and only if the intermediate double cover is smooth. The intermediate double cover has equation $z^2 - 2ag^nz + g^n = 0$ and its branch divisor is $\{g^n(g^na^2 - 1) = 0\} = \{a^2 - F^n = 0\}$. By hypothesis (ii) this locus is smooth, so X is smooth where $v_1u_1 \neq 0$.

It remains to consider the case where $v_1u_0 \neq 0$. Here, setting $v_1 = 1 = u_0$, the equations (31) reduce to

$$\begin{cases} u_1 - v_0F & = 0 \\ a_\infty - 2a_0v_0^n + a_\infty v_0^n u_1^n & = 0. \end{cases}$$

Substituting $u_1 = v_0F$ in the second equation we obtain:

$$a_\infty - 2a_0v_0^n + a_\infty F^n v_0^{2n} = 0.$$

Notice that, if $v_0 \neq 0$, then we have already seen that X is smooth. On the other hand, when $v_0 = 0$, we must have $a_\infty = 0$, and then the smoothness of A_∞ implies that of X .

The finiteness of π follows from the fact that π has finite fibre and its restriction to each open subset of Y where \mathbb{L} is trivial is projective ([Hart77], III, Ex. 11.1).

To describe the branch divisor, recall that the restriction of π to the open set $u_0v_0 \neq 0$ is a simple dihedral cover and so its branch divisor is $F^n - a^2 = 0$, where $a := a_0/a_\infty$. Notice that, if $u_0v_0 \neq 0$, then $a_\infty \neq 0$ everywhere, on the other hand, if $u_0v_0 = 0$, then $u_1v_1 = 0$ by (31), so $X \cap \{u_0v_0 = 0\} \subset \{u_1v_0 \neq 0\} \cup \{u_0v_1 \neq 0\}$. The claim now follows from the previous explicit description of π on $u_0v_1 \neq 0$.

Finally, if $A_0 \cap \{F = 0\} \neq \emptyset$, then X is irreducible since the open subset $X \cap \{u_0v_0 \neq 0\}$ is irreducible by Theorem 6.1. \square

The invariants of almost simple D_n -covers $\pi: X \rightarrow Y$ can be computed in the same way as in the simple case, once a description of $\pi_*\mathcal{O}_X$ in terms of L and A_∞ is provided. In the remaining part of this section we show that there is an isomorphism as follows:

$$(32) \quad \pi_*\mathcal{O}_X \cong \mathcal{O}_Y \oplus \mathcal{O}_Y(-nL - A_\infty) \oplus \left(\bigoplus_{i=1}^{n-1} [\mathcal{O}_Y(-iL) \oplus \mathcal{O}_Y(-(n-i)L)] \right) (-A_\infty).$$

Notice that, on the open subset $Y \setminus A_\infty$ where π is a simple cover, the previous formula reduces to (27).

To prove (32), we consider the following Cartesian diagram

$$\begin{array}{ccc} \mathcal{Q} & \xrightarrow{\tilde{p}_2} & \mathbb{P}_2 \\ \tilde{p}_1 \downarrow & & \downarrow p_2 \\ \mathbb{P}_1 & \xrightarrow{p_1} & Y \end{array}$$

where $\mathcal{Q} := \mathbb{P}_1 \times_Y \mathbb{P}_2$ is a $\mathbb{P}^1 \times \mathbb{P}^1$ -bundle with projection $p: \mathcal{Q} \rightarrow Y$, $p = p_i \circ \tilde{p}_i$, $\forall i = 1, 2$. Recall that the Picard group of \mathcal{Q} is isomorphic

to $\text{Pic}(Y) \times \mathbb{Z}^{\oplus 2}$ via the usual isomorphism that sends $(\mathcal{L}, m, n) \in \text{Pic}(Y) \times \mathbb{Z}^{\oplus 2}$ to $p^*\mathcal{L} \otimes \mathcal{O}_{\mathcal{Q}}(m, n) := p^*\mathcal{L} \otimes \mathcal{O}_{\mathbb{P}_1}(m) \boxtimes \mathcal{O}_{\mathbb{P}_2}(n)$.

Let us define $D_i := \{\Phi_i = 0\}$ to be the divisor of \mathcal{Q} given by the equation $\Phi_i = 0$ in (31), for $i = 1, 2$, and notice that

$$(33) \quad \begin{aligned} \Phi_1 &\in H^0(\mathcal{Q}, \mathcal{O}_{\mathcal{Q}}(1, 1) \otimes p^*\mathcal{O}_Y(2L)) , \\ \Phi_2 &\in H^0(\mathcal{Q}, \mathcal{O}_{\mathcal{Q}}(n, n) \otimes p^*\mathcal{O}_Y(nL + A_{\infty})) . \end{aligned}$$

Then we consider the usual short exact sequence

$$(34) \quad 0 \rightarrow \mathcal{I}_X \rightarrow \mathcal{O}_{\mathcal{Q}} \rightarrow \mathcal{O}_X \rightarrow 0 ,$$

where \mathcal{I}_X is the sheaf of ideals of X . Since X is the complete intersection of two divisors in \mathcal{Q} , the Koszul resolution of \mathcal{I}_X is as follows:

$$(35) \quad 0 \rightarrow \mathcal{O}_{\mathcal{Q}}(-D_1 - D_2) \rightarrow \mathcal{O}_{\mathcal{Q}}(-D_1) \oplus \mathcal{O}_{\mathcal{Q}}(-D_2) \rightarrow \mathcal{I}_X \rightarrow 0 .$$

Applying p_* to (34) we obtain the following split short exact sequence:

$$0 \rightarrow \mathcal{O}_Y \cong p_*\mathcal{O}_{\mathcal{Q}} \rightarrow p_*\mathcal{O}_X \rightarrow R^1p_*\mathcal{I}_X \rightarrow 0 ,$$

where we have used the fact that $p_*\mathcal{I}_X = 0 = R^1p_*\mathcal{O}_{\mathcal{Q}}$.

In order to compute $R^1p_*\mathcal{I}_X$, we apply p_* to (35) and we obtain the following exact sequence:

$$0 \rightarrow R^1p_*\mathcal{I}_X \rightarrow R^2p_*\mathcal{O}_{\mathcal{Q}}(-D_1 - D_2) \rightarrow R^2p_*(\mathcal{O}_{\mathcal{Q}}(-D_1) \oplus \mathcal{O}_{\mathcal{Q}}(-D_2)) ,$$

where we have used the equality $R^1p_*(\mathcal{O}_{\mathcal{Q}}(-D_1) \oplus \mathcal{O}_{\mathcal{Q}}(-D_2)) = 0$ that follows from the Künneth formula. Furthermore, since $R^2p_*\mathcal{O}_{\mathcal{Q}}(-D_1) = 0$, we have that

$$R^1p_*\mathcal{I}_X = \ker [R^2p_*\mathcal{O}_{\mathcal{Q}}(-D_1 - D_2) \xrightarrow{\mu} R^2p_*\mathcal{O}_{\mathcal{Q}}(-D_2)] ,$$

where μ is induced by the morphism $\mathcal{O}_{\mathcal{Q}}(-D_1 - D_2) \rightarrow \mathcal{O}_{\mathcal{Q}}(-D_2)$ in the Koszul resolution of \mathcal{I}_X , given by $\psi_1 \wedge \psi_2 \mapsto \psi_1(\Phi_1)\psi_2$, for $\psi_i \in \mathcal{O}_{\mathcal{Q}}(-D_i)$.

To describe μ explicitly, we use the following isomorphisms:

$$\begin{aligned} R^2p_*\mathcal{O}_{\mathcal{Q}}(-D_1 - D_2) &\cong S^{n-1}(\mathcal{O}_Y \oplus \mathcal{O}_Y(L))^{\otimes 2} \otimes \mathcal{O}_Y(-nL - A_{\infty}) , \\ R^2p_*\mathcal{O}_{\mathcal{Q}}(-D_2) &\cong S^{n-2}(\mathcal{O}_Y \oplus \mathcal{O}_Y(L))^{\otimes 2} \otimes \mathcal{O}_Y(-(n-2)L - A_{\infty}) , \end{aligned}$$

that follow applying the projection formula, Künneth formula, and the standard isomorphisms (see e.g. [Hart77, Ex. 8.4, III]). Then, if we choose local sections x_0, y_0 of \mathcal{O}_Y , and x_1, y_1 of $\mathcal{O}_Y(L)$, that generate the corresponding sheaves, we obtain the following local basis for $S^{n-1}(\mathcal{O}_Y \oplus \mathcal{O}_Y(L))^{\otimes 2}$:

$$E_{ij} := x_0^i x_1^{n-1-i} \otimes y_0^j y_1^{n-1-j} , \quad 0 \leq i, j \leq n-1 .$$

Similarly, $G_{km} := (x_0^k x_1^{n-2-k} \otimes y_0^m y_1^{n-2-m}) \otimes (x_1 \otimes y_1)$, for $0 \leq k, m \leq n-2$, is a local basis for $S^{n-2}(\mathcal{O}_Y \oplus \mathcal{O}_Y(L))^{\otimes 2} \otimes \mathcal{O}_Y(2L)$. The morphism μ in these basis is given as follows:

$$\mu(E_{ij}) = G_{ij} - fG_{i-1, j-1} ,$$

where $F = fx_1 \otimes y_1$, and $G_{km} := 0$, for $k, m \notin \{0, \dots, n-2\}$. Hence from elementary linear algebra, we have that $\ker(\mu)$ (twisted by $\mathcal{O}_Y(nL + A_\infty)$) is generated by $E_{n-1,j}$ and $E_{i,n-1}$, for $0 \leq i, j \leq n-1$. So

$$\ker(\mu) = \left([\oplus_{j=0}^{n-1} \mathcal{O}_Y((n-1-j)L)] \oplus [\oplus_{i=0}^{n-2} \mathcal{O}_Y((n-1-i)L)] \right) \otimes \mathcal{O}_Y(-nL - A_\infty),$$

and hence (32) follows.

7. DEFORMATIONS OF SIMPLE DIHEDRAL COVERS

Let X be a simple dihedral covering of Y : this means that X is the subvariety of the vector bundle $V = \mathbb{L} \oplus \mathbb{L}$ which is (see 26) the complete intersection of two hypersurfaces, one in $|p^*(2L)|$, the other in $|p^*(nL)|$; here $p: V \rightarrow Y$ is the natural projection.

Observe now that the cotangent sheaf of V is an extension of $p^*\Omega_Y^1$ by $p^*(\mathcal{O}_Y(-L)^{\oplus 2})$,

$$(36) \quad 0 \rightarrow p^*\Omega_Y^1 \rightarrow \Omega_V^1 \rightarrow p^*(\mathcal{O}_Y(-L)^{\oplus 2}) \rightarrow 0,$$

where $p^*\Omega_Y^1 \rightarrow \Omega_V^1$ is the cotangent map of p .

Then the conormal sheaf exact sequence of X reads out, if we denote by $L' := \pi^*(L)$, as

$$0 \rightarrow N_{X|V}^* = \mathcal{O}_X(-2L') \oplus \mathcal{O}_X(-nL') \rightarrow \Omega_V^1 \otimes \mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow 0,$$

and the dual sequence is

$$0 \rightarrow \Theta_X \rightarrow \Theta_V \otimes \mathcal{O}_X \rightarrow N_{X|V} = \mathcal{O}_X(2L') \oplus \mathcal{O}_X(nL') \rightarrow 0,$$

whose direct image under π_* yields the tangent exact sequence:

$$0 \rightarrow \pi_*\Theta_X \rightarrow \pi_*(\Theta_V \otimes \mathcal{O}_X) \rightarrow (\mathcal{O}_Y(2L) \oplus \mathcal{O}_Y(nL)) \otimes \pi_*(\mathcal{O}_X) \rightarrow 0.$$

Passing to the long exact cohomology sequence, we get the Kodaira-Spencer exact sequence

$$(37) \quad H^0((\mathcal{O}_Y(2L) \oplus \mathcal{O}_Y(nL)) \otimes \pi_*(\mathcal{O}_X)) \rightarrow H^1(\Theta_X) \rightarrow H^1(\pi_*(\Theta_V \otimes \mathcal{O}_X)) \rightarrow \dots$$

The meaning of the first linear map is given through the following definition.

Definition 7.1. *The space of natural deformations of a simple dihedral covering is the family of complete intersections of V :*

$$(38) \quad \begin{cases} uv - F & = 0 \\ u^n - 2a + v^n + \sum_1^{n-1} (b_i u^i + c_i v^i) + d(u^n - v^n) & = 0, \end{cases}$$

$$(39) \quad \text{where } b_i, c_i \in H^0(\mathcal{O}_Y(n-i)L), d \in H^0(\mathcal{O}_Y).$$

Remark 7.2. The reader can see that in the second equation

$$-2a + v^n + \sum_1^{n-1} (b_i u^i + c_i v^i) + d(u^n - v^n)$$

can be any section in $H^0(\mathcal{O}_Y(nL) \otimes \pi_*(\mathcal{O}_X))$, in view of our basic formulae; instead, any section in $H^0(\mathcal{O}_Y(2L) \otimes \pi_*(\mathcal{O}_X))$ is of the form

$$-F + \beta u + \alpha v + \lambda u^2 + \mu v^2, \text{ where } \alpha, \beta \in H^0(\mathcal{O}_Y(L)), \lambda, \mu \in \mathbb{C}.$$

But the new equation $uv - F + \beta u + \alpha v + \lambda v^2 + \mu u^2$ is the old form $u'v' = F'$ (up to a multiplicative constant) if we choose new variables

$$u' := u + \alpha + \lambda v, \quad v' := v + \beta + \mu u.$$

Of course, one can deform not only the equations, but also simultaneously the base Y , the vector bundle V , and the equations; this however leads in general to a deformation with non smooth base.

We have at any rate an easy result which says that all small deformations are obtained by natural deformations.

Theorem 7.3. *Assume that $\pi: X \rightarrow Y$ is a simple dihedral covering. Then all small deformations of X are natural deformations of $\pi: X \rightarrow Y$, provided $H^1(\pi_*(\Theta_V \otimes \mathcal{O}_X)) = 0$ (that happens, for example if $H^1((\Theta_Y \oplus \mathcal{O}_Y(L)^{\oplus 2}) \otimes \pi_*(\mathcal{O}_X)) = 0$). In particular, the Kuranishi family of X is smooth, and the Kuranishi space $\text{Def}(X)$ is locally analytically isomorphic to*

$$\text{Def}' := \text{coker} \left(H^0(\pi_*(\Theta_V \otimes \mathcal{O}_X)) \rightarrow H^0((\mathcal{O}_Y(2L) \oplus \mathcal{O}_Y(nL)) \otimes \pi_*(\mathcal{O}_X)) \right).$$

Proof. By our assumption, and the Kodaira-Spencer exact sequence, we have an isomorphism of vector spaces $\text{Def}' \cong H^1(\Theta_X)$. Moreover, by the previous remark, the family of natural deformations has Kodaira-Spencer map which is surjective onto Def ; therefore, by the implicit functions theorem, $\text{Def}(X)$ is the germ of the analytic space $H^1(\Theta_X)$ at the origin, hence our claim. \square

8. EXAMPLES AND APPLICATIONS

8.1. Simple dihedral covers of projective spaces. We first consider the case where $Y = \mathbb{P}^2$, the complex projective plane, $n = 3$ and $\mathcal{O}_Y(L) = \mathcal{O}_{\mathbb{P}^2}(1)$. Then $a \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(3))$ and $F \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2))$ are a cubic and a quadric curve respectively, such that the sextic curve $\{F^3 - a^2 = 0\}$ is smooth in the locus $F \neq 0$, and $\{a = 0\}$ intersects $\{F = 0\}$ transversely. By Theorem 6.1, we have a smooth D_3 -cover $\pi: X \rightarrow \mathbb{P}^2$ branched over $\mathcal{B} = \{F^3 - a^2 = 0\}$. Notice that $\mathcal{B} \in |-2K_{\mathbb{P}^2}|$ and $\omega_{\mathbb{P}^2} = \mathcal{O}_Y(-3L)$. Hence

$$\omega_X = \pi^*(\omega_{\mathbb{P}^2}(3L)) \cong \mathcal{O}_X.$$

Furthermore, $q(X) = 0$, hence X is a K3 surface.

Let $W = X/\langle \tau \rangle$ and let $f: W \rightarrow \mathbb{P}^2$ be the induced triple cover (see Section 5.3). Then W can be realised as a cubic surface in \mathbb{P}^3 in such a

way that f is the projection from a point in $\mathbb{P}^3 \setminus W$. Indeed, consider the equations (26) which define X . If we define $w := u + v$, then

$$\begin{aligned} w^3 &= u^3 + v^3 + 3uv(u + v) \\ &= 2a + 3Fw. \end{aligned}$$

The branch divisor of f is $\mathcal{B} = \{F^3 - a^2\}$. So, under the hypotheses of Theorem 6.1, \mathcal{B} is a sextic with 6 cusps lying on a conic.

The fundamental group of the complement $\mathbb{P}^2 \setminus \mathcal{B}$ of a sextic curve \mathcal{B} as above has been studied in [Zar29], where in particular it is proven that $\pi_1(\mathbb{P}^2 \setminus \mathcal{B})$ is generated by two elements of order 2 and 3 respectively. From this it follows that there exists a surjective group homomorphism $\pi_1(\mathbb{P}^2 \setminus \mathcal{B}) \rightarrow D_3$, and hence it follows from the generalised Riemann existence theorem of Grauert and Remmert, that there exists a D_3 -cover $\pi: X \rightarrow \mathbb{P}^2$. From Theorem 6.1 we have an explicit construction of such a cover.

Let now $n > 3$ and consider the simple D_n -cover $\pi: X \rightarrow \mathbb{P}^2$ associated to $\mathcal{O}_Y(L) = \mathcal{O}_{\mathbb{P}^2}(1)$, $a \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(n))$ and $F \in H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(2))$. Under the hypotheses of Theorem 6.1, X is a smooth surface with

$$\begin{aligned} \omega_X &= \pi^* \mathcal{O}_{\mathbb{P}^2}(n - 3), \\ K_X^2 &= 2n(n - 3)^2, \\ \chi(\mathcal{O}_X) &= \frac{1}{3}n^3 - \frac{3}{2}n^2 + \frac{13}{6}n, \end{aligned}$$

in particular X is a surface of general type, which is minimal since $K_X \cdot C > 0$ for any curve $C \subset X$.

Finally, let $n = 2$, $Y = \mathbb{P}^2$ and $\mathcal{O}_Y(L) = \mathcal{O}_{\mathbb{P}^2}(1)$. Under the hypotheses of Theorem 6.1, X is a smooth surface with the following invariants:

$$\begin{aligned} \omega_X &= \pi^* \mathcal{O}_{\mathbb{P}^2}(-1) \\ \chi(\mathcal{O}_X) &= 1 \\ q &= 0 \\ K_X^2 &= 4 \\ p_n &= \dim H^0(X, \omega_X^{\otimes n}) = 0. \end{aligned}$$

Hence X is a rational, non-minimal surface. Indeed, X is isomorphic to a del Pezzo surface of degree 4 in \mathbb{P}^4 , the complete intersection of the quadrics $uv = F(x_0 : x_1 : x_2)$, $u^2 + v^2 = 2a(x_0 : x_1 : x_2)$, where $(x_0 : x_1 : x_2 : u : v)$ are now homogeneous coordinates in \mathbb{P}^4 .

In a similar way one can construct examples of dihedral covers of \mathbb{P}^2 with $\mathcal{O}_Y(L) = \mathcal{O}_{\mathbb{P}^2}(d)$, $d > 1$.

As an application of Theorem 7.3, we have that all the small deformations of a simple D_n -cover $\pi: X \rightarrow \mathbb{P}^2$ associated to $\mathcal{O}_Y(L) = \mathcal{O}(m)$, $m \geq 1$, are natural deformations, if $(m, n) = (1, 2)$, or $m \geq 2$ and any

$n \geq 2$. This follows directly from Thm. 7.3, formula (27) and the computation of the cohomology of $\Omega_{\mathbb{P}^d}^q(k)$ ([Bott57, p. 256]).

Remark 8.1. Notice that, in general, a simple D_n -cover of \mathbb{P}^d , $\pi: X \rightarrow \mathbb{P}^d$, associated to $\mathcal{O}_Y(L) = \mathcal{O}(m)$, $F \in H^0(\mathbb{P}^d, \mathcal{O}(2m))$ and $a \in H^0(\mathbb{P}^d, \mathcal{O}(nm))$, is isomorphic to a complete intersection X' in the weighted projective space $\mathbb{P}^{d+2}(1, \dots, 1, m, m)$, where

$$X' := \{(x_0 : \dots : x_d : u : v) \in \mathbb{P}^{d+2}(1, \dots, 1, m, m) \mid uv = F, u^n + v^n = 2a\}.$$

To see this, observe that $X \subset \mathbb{L} \oplus \mathbb{L}$ is the quotient of $\tilde{X} \subset (\mathbb{C}^{d+1} \setminus \{0\}) \times \mathbb{C}^2$ via the linear diagonal action of \mathbb{C}^* with weights $(1, \dots, 1, m, m)$, where

$$\tilde{X} := \{(x_0, \dots, x_d, u, v) \mid uv = F, u^n + v^n = 2a\}.$$

The claim now follows since $(\mathbb{C}^{d+1} \setminus \{0\}) \times \mathbb{C}^2 \subset \mathbb{C}^{d+3} \setminus \{0\}$, and the action of \mathbb{C}^* on $(\mathbb{C}^{d+1} \setminus \{0\}) \times \mathbb{C}^2$ is the restriction of the linear diagonal action on $\mathbb{C}^{d+3} \setminus \{0\}$ with weights $(1, \dots, 1, m, m)$.

If $d \geq 3$, then Theorem 7.3 implies that all the small deformations of X are natural deformations of the simple D_n -cover $\pi: X \rightarrow \mathbb{P}^d$. Indeed, in this case, we have that

$$H^1((\Theta_{\mathbb{P}^d} \oplus \mathcal{O}_{\mathbb{P}^d}(m))^{\oplus 2} \otimes \pi_* \mathcal{O}_X) = 0,$$

as it follows from the fact that $H^1(\mathbb{P}^d, \mathcal{O}_{\mathbb{P}^d}(k)) = 0$, $\forall k$, and that $H^1(\mathbb{P}^d, \Theta_{\mathbb{P}^d}(k)) \cong H^{d-1}(\mathbb{P}^d, \Omega_{\mathbb{P}^d}^1(-k-d-1))^\vee = 0$, $\forall k$ ([Bott57]).

8.2. An application to fundamental groups. According to the generalized Riemann existence theorem of Grauert and Remmert [GR58], coverings $\pi: X \rightarrow Y$ of a normal variety Y , of degree n , with branch locus contained in a divisor $\mathcal{B} \subset Y$, and with X normal, correspond to conjugacy classes of group homomorphisms $\mu: \pi_1(Y \setminus \mathcal{B}) \rightarrow \mathfrak{S}_n$. In this situation, X is irreducible, if and only if $\text{Im}(\mu)$ is a transitive subgroup of \mathfrak{S}_n ; $\pi: X \rightarrow Y$ is Galois with group $G := \text{Im}(\mu)$, if and only if G coincides with the group of automorphisms of π . In particular, if $\pi: X \rightarrow Y$ is a G -cover with G non-abelian and X irreducible, then $\pi_1(Y \setminus \mathcal{B})$ is necessarily non-abelian.

Fundamental groups of complements of divisors in projective varieties have been extensively studied by many authors. As a direct consequence of Theorem 6.1, we have the following result.

Proposition 8.2. *Let Y be a smooth variety and $L \subset Y$ be a divisor. Assume that there exist $a \in H^0(Y, \mathcal{O}_Y(nL))$ and $F \in H^0(Y, \mathcal{O}_Y(2L))$, such that the conditions (i), (ii) of Thm. 6.1 are satisfied and $\{a = 0\} \cap \{F = 0\} \neq \emptyset$. Then $\pi_1(Y \setminus \mathcal{B})$ admits an epimorphism onto D_n , in particular it is non-abelian, where $\mathcal{B} = \{a^2 - F^n = 0\}$.*

Notice that similar results have been obtained using different methods ([CKO03, Lemma 3], [ArCo10]). Briefly, one considers the pencil $\{\lambda 2(a = 0) + \mu n(F = 0)\}_{(\lambda: \mu) \in \mathbb{P}^1} \subset |2nL|$ and the induced morphism $Y \setminus \mathcal{B} \rightarrow \mathbb{P}^1 \setminus \{(1 : 1)\}$. This gives a group homomorphism

$\pi_1(Y \setminus \mathcal{B}) \rightarrow \pi_1^{orb}(\mathbb{P}^1 \setminus \{(1 : 1)\})$, where π_1^{orb} is the orbifold fundamental group of $\mathbb{P}^1 \setminus \{(1 : 1)\}$ with two orbifold points, $(1 : 0)$ of order 2 and $(0 : 1)$ of order n . Now, using the long exact homotopy sequence, one concludes that $\pi_1(Y \setminus \mathcal{B}) \rightarrow \pi_1^{orb}(\mathbb{P}^1 \setminus \{(1 : 1)\})$ is surjective and so $\pi_1(Y \setminus \mathcal{B})$ is not abelian.

REFERENCES

- [ArCo10] Artal-Bartolo, E.; Cogolludo-Agustín, J.; *On the connection between fundamental groups and pencils with multiple fibers.* J. Singul. 2 (2010), 1–18.
- [BaCa08] Bauer, I.; Catanese, F. *A volume maximizing canonical surface in 3-space.* Comment. Math. Helv. 83 (2008), no. 2, 387–406.
- [Bott57] Bott, R. *Homogeneous vector bundles.* Ann. of Math. (2) 66 (1957), 203–248.
- [Bur66] Burniat, P., *Sur les surfaces de genre $P_{12} > 1$.* Ann. Mat. Pura Appl. (4) 71, 1–24 (1966).
- [Cat84] Catanese, F. *On the moduli spaces of surfaces of general type.* J. Differential Geom. 19 (1984), no. 2, 483–515.
- [Cat88] Catanese, F. *Moduli of algebraic surfaces.* Theory of moduli (Montecatini Terme, 1985), 1–83, Lecture Notes in Math., 1337, Springer, Berlin, (1988).
- [Cat89] Catanese, F., *Everywhere non reduced moduli spaces.* Invent. math. 98, 293–310 (1989).
- [Cat98] Catanese, F., *Singular bidouble covers and the construction of interesting algebraic surfaces.* Algebraic geometry: Hirzebruch 70 (Warsaw, 1998), 97–120, Contemp. Math., 241, Amer. Math. Soc., Providence, RI, 1999.
- [CKO03] Catanese, F.; Keum, J.; Oguiso, K.; *Some remarks on the universal cover of an open K3 surface.* Math. Ann. 325 (2003), no. 2, 279–286.
- [Cat10] Catanese, F., *Irreducibility of the space of cyclic covers of algebraic curves of fixed numerical type and the irreducible components of $Sing(\overline{\mathfrak{M}}_g)$.* Advances in geometric analysis, 281–306, Adv. Lect. Math. (ALM), 21, Int. Press, Somerville, MA, 2012.
- [CHKS06] Catanese, F.; Hoşten, S.; Khetan, A.; Sturmfels, B., *The maximum likelihood degree.* Amer. J. Math. 128 (2006), no. 3, 671–697.
- [CLP] Catanese, F., Lönne, M., Perroni, F. (In preparation.)
- [Com30] Comessatti, A. *Sulle superficie multiple cicliche.* Rendiconti Seminario Padova 1, 1–45 (1930)
- [East11] Easton, R. W., *S_3 -covers of schemes.* Canad. J. Math. 63 (2011), no. 5, 1058–1082.
- [Enr49] Enriques, F., *Le Superficie Algebriche.* Nicola Zanichelli, Bologna (1949).
- [GH] Griffiths, P.; Harris, J.; Principles of algebraic geometry. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994.
- [GR58] Grauert, H.; Remmert, R.; *Komplexe Räume.* Math. Ann. 136 (1958), 245–318.
- [Hart77] Hartshorne, R., Algebraic geometry. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977.
- [Mat89] Matsumura, H. Commutative ring theory. Second edition. Cambridge Studies in Advanced Mathematics, 8. Cambridge University Press, Cambridge, 1989.

- [Mir85] Miranda, R. *Triple covers in algebraic geometry*. Amer. J. Math. 107 (1985), no. 5, 1123–1158.
- [Mum84] Mumford, D., *Tata lectures on theta. II*. Reprint of the 1984 original. Birkhäuser Boston, Inc., Boston, MA, 2007.
- [Namba87] Namba, M. *Branched coverings and algebraic functions*. Pitman Research Notes in Mathematics Series, 161. Longman Scientific & Technical, Harlow; John Wiley & Sons, Inc., New York, 1987.
- [Par91] Pardini, R. *Abelian covers of algebraic varieties*. J. Reine Angew. Math. 417 (1991), 191–213.
- [ReidC3f] Reid, M., *Canonical 3-folds*. Journées de Géométrie Algébrique d'Angers, Juillet 1979/Algebraic Geometry, Angers, 1979, pp. 273–310, Sijthoff & Noordhoff, Alphen aan den Rijn–Germantown, Md., 1980.
- [Sai80] Saito, K., *Theory of logarithmic differential forms and logarithmic vector fields*. J. Fac. Sci. Univ. Tokyo Sect. IA Math. 27 (1980), no. 2, 265–291.
- [Serre77] Serre, J.-P., *Linear representations of finite groups*. Graduate Texts in Mathematics, Vol. 42. Springer-Verlag, New York-Heidelberg, 1977.
- [Tok94] Tokunaga, H.-O., *On dihedral Galois coverings*. Canad. J. Math. 46 (1994), no. 6, 1299–1317.
- [Zar29] Zariski, O., *On the Problem of Existence of Algebraic Functions of Two Variables Possessing a Given Branch Curve*. Amer. J. Math. 51 (1929), no. 2, 305–328.
- [Zar58] Zariski, O., *On the purity of the branch locus of algebraic functions*. Proc. Nat. Acad. Sci. U.S.A. 44 1958 791–796.

Authors' Address:

Fabrizio Catanese,
Lehrstuhl Mathematik VIII,
Mathematisches Institut der Universität Bayreuth
NW II, Universitätsstr. 30
95447 Bayreuth (Germany).

Fabio Perroni,
Dipartimento di Matematica e Geoscienze,
Sezione di Matematica e Informatica, Università di Trieste, Via Valerio
12/1, 34127 Trieste (Italy).

Emails: Fabrizio.Catanese@uni-bayreuth.de; fperroni@units.it.